

A generalization of the Jiang-Su construction

Yasuhiko Sato

Department of Mathematics, Hokkaido University,
Sapporo 060-0810, Japan
e-mail : s053018@math.sci.hokudai.ac.jp

Abstract

Let G be a countable abelian group. We construct a unital simple projectionless C^* -algebra A with a unique tracial state, that satisfies $(K_0(A), [1_A]) \cong (\mathbb{Z}, 1)$, $K_1(A) \cong G$, $A \otimes \mathcal{Z} \cong A$, and that is obtained as the inductive limit C^* -algebra of a sequence of dimension drop algebras of a specific form. This construction is based on the construction of the Jiang-Su algebra. By this construction, we show a certain conjugacy result for aperiodic automorphisms of these projectionless C^* -algebras. We also show that an automorphism of this projectoinless C^* -algebra has a certain aperiodicity up to the weakly inner automorphisms in the tracial representation if and only if it has a kind of Rohlin property, which leads to the Rohlin property after taking tensor product of certain C^* -algebras of real rank zero.

2000 Mathematics Subject Classification. Primary 46L35; Secondary 46L80, 46L40.

Key words and phrases. C^* -algebra, K-theory, Jiang-Su algebra, automorphism, Rohlin property.

1 Introduction

In [6], Jiang and Su have concretely constructed the projectionless C^* -algebra $\mathcal{Z}$ whose K_0 -group is $(\mathbb{Z}, \mathbb{Z}_+, 1)$ and K_1 -group is $\{0\}$ which is now called the Jiang-Su algebra and they have shown that any automorphism of the Jiang-Su algebra is approximately inner. For the classification program of finite C^* -algebras started by G. A. Elliott, the Jiang-Su algebra behaves as $\mathcal{O}_\infty$ in the classification theory for Kirchberg algebras. Then the Jiang-Su algebra was studied by many people (e.g., [2], [3], [12], [15], and [16]).

In this paper elaborating Jiang-Su's method we give a unified way of constructing unital projectionless C^* -algebras with arbitrary K_1 -group. And, as an application of this construction, we can prove the following corollary as remarked in Remark 5.3.

Corollary 1.1. *Let A be the unital simple projectionless C^* -algebra that is obtained in Theorem 3.2 or 4.5, and τ the unique tracial state of A . Suppose that α and β are automorphisms of A which are aperiodic in the quotient*

$\text{Aut}(A)/\text{WInn}(A)$, then for any $\varepsilon > 0$ there exists an approximately inner automorphism $\sigma \in \text{Aut}(A)$ and $\gamma \in \text{WInn}(A)$ with $W \in U(\pi_\tau(A)'')$ such that $\pi_\tau \circ \gamma = \text{Ad } W \circ \pi_\tau$, $\|W - 1\|_2 < \varepsilon$, and

$$\alpha = \gamma \circ \sigma \circ \beta \circ \sigma^{-1}.$$

Here $\text{WInn}(A)$ is the subgroup of the automorphism group of A defined by

$$\text{WInn}(A) = \{\alpha \in \text{Aut}(A); \pi_\tau \circ \alpha = \text{Ad } V \circ \pi_\tau, V \in U(\pi_\tau(A)'')\},$$

and π_τ is the GNS-representation associated with τ . This corollary is a straightforward generalization of Theorem 2.6 in [13]. The aperiodicity in the assumption of this corollary will be characterized by the following theorem.

Theorem 1.2. *Let A and τ be as in the above corollary and α is an automorphism of A . Then $[\alpha] \in \text{Aut}(A)/\text{WInn}(A)$ is aperiodic if and only if α satisfies the following condition that: for any $k \in \mathbb{N}$ there exists a central sequence $(f_n)_n$ of positive elements in A (i.e., $(f_n)_n \in (A_\infty)_+^1$) such that*

$$(f_n)_n \cdot \alpha^j((f_n)_n) = 0, \quad j = 1, 2, \dots, k-1, \quad \tau(1_A - \sum_{j=0}^{k-1} \alpha^j(f_n)) \rightarrow 0,$$

This theorem is motivated by a similar result for UHF-algebras by Kishimoto [7]. The condition expressed in terms of positive elements in this theorem, has given the result in [13] that: when A is the Jiang-Su algebra, B is a unital simple TAF-algebra satisfying a certain condition in [13], β is an automorphism of B , and $\alpha \in \text{Aut}(A)$ has the property in the above theorem, this condition implies that the automorphism $\alpha \otimes \beta$ of $A \otimes B$ is asymptotically unitarily equivalent to an automorphism with the Rohlin property.

The building blocks of the unital simple projectionless C^* -algebras constructed in Theorem 3.2 or 4.5 will be dimension drop algebras on one-dimensional compact Hausdorff graphs with ‘dimension drop’ condition at each end point. Give a finitely generated abelian group G , the base space is uniquely determined as a one-point union of copies of torus $\mathbb{T}$ and copies of the interval $[0, 2\pi]$ with $0 \in [0, 2\pi]$ the common point, where the number of copies of $\mathbb{T}$ corresponds to the rank of the free part of G and the number of copies of $[0, 2\pi]$ corresponds to the number of directly indecomposable summands of the torsion subgroup of G . Regulating matrix algebras on the end points $2\pi \in [0, 2\pi]$, we obtain a unital projectionless dimension drop algebra that has the finite cyclic groups as K_1 -group. In [4], Elliott have already obtained the projectionless C^* -algebra whose K_0 -group is $(\mathbb{Z}, \mathbb{Z}_+, 1)$ and these K_1 -group is a given countable abelian group, and many people have studied subhomogeneous C^* -algebras on one dimensional compact Hausdorff spaces of similar type (see e.g. [8], [10], [14]). However the author could not find a type of unital projectionless dimension drop algebras suitable to prove Corollary 1.1. Recently, H. Lin showed the classification result of the inductive limit C^* -algebras of dimension drop algebras of general type, in [9].

The contents of this paper are as follows. In Section 2, for a given finitely generated abelian group G we construct a projectionless dimension drop algebra whose K_1 -group is G . In Section 3, for a given countable abelian group G we construct a unital simple projectionless C^* -algebra whose K_1 -group is G that is the inductive limit C^* -algebra of a sequence of dimension drop algebras. In Section 4, we show $\mathcal{Z}$ -stability of these projectionless C^* -algebras, which is a key property in the classification theory according to [5] and [16]. In Section 5, we prove Corollary 1.1 and Theorem 1.2. In order to show this corollary we need to show a UHF-embeddability, and in order to show this UHF-embeddability we also need to consider another one-dimensional homogeneous C^* -algebra which naturally contains the dimension drop algebra mentioned above.

Concluding this section, we prepare some notations. When A is a C^* -algebra, we denote by A^1 the set $\{a \in A : \|a\| \leq 1\}$, A_{sa} the set of self-adjoint elements of A , $U(A)$ the unitary group of A , $T(A)$ the tracial state space of A , A^∞ the quotient $\ell^\infty(\mathbb{N}, A)/c_0(A)$, and A_∞ the relative commutant $A^\infty \cap A'$. We use M_n to denote the C^* -algebra of $n \times n$ matrices with complex entries and we denote by Tr_n the normalized trace on M_n . In this paper we denote by p_i the i th-prime number and (m, n) the greatest common divisor of m and n . On a metric space (X, d) , we say that $Y \subset X$ is ε -dense in X if for $y \in Y$ there is $x \in X$ such that $d(x, y) \leq \varepsilon$.

2 Dimension drop algebras

First we shall prepare some symbols and define a dimension drop algebra for a finitely generated abelian group. For a given countable abelian group G we will express G as the inductive limit of an increasing sequence of finitely generated abelian group G_n , $n \in \mathbb{N}$, and construct a unital simple projectionless C^* -algebra whose K_1 -group is G as the inductive limit C^* -algebra of dimension drop algebras for G_n , $n \in \mathbb{N}$. Then we denote by G_n a finitely generated abelian group, by $G_{n,0}$ the free part of G_n , and by $G_{n,1}$ the torsion subgroup of G_n . Set $r_0^{(n)} = \text{rank}(G_{n,0}) \in \mathbb{Z}_+$. For $i \in \mathbb{N}$ with $p_i \mid |G_{n,1}|$, let $G_{n,i}$ be the subgroup of G_n generated by $\{g \in G_n; o(g) = p_i^n, n \in \mathbb{N}\}$, where $o(g)$ means the order of g , and let $r_i^{(n)} \in \mathbb{N}$ and $d_{i,j}^{(n)} \in \mathbb{N}$, $j = 1, 2, \dots, r_i^{(n)}$ be such that

$$G_{n,i} \cong \bigoplus_{j=1}^{r_i^{(n)}} \mathbb{Z}_{d_{i,j}^{(n)}}, \quad d_{i,j}^{(n)} = p_i^{k_{i,j}^{(n)}},$$

where $\mathbb{Z}_{d_{i,j}^{(n)}} = \mathbb{Z}/d_{i,j}^{(n)}\mathbb{Z}$, $k_{i,j}^{(n)} \in \mathbb{N}$ and $k_{i,j}^{(n)} \leq k_{i,j+1}^{(n)}$. Therefore we identify

$$G_n \cong G_{n,0} \oplus G_{n,1} \cong \mathbb{Z}^{r_0^{(n)}} \oplus \bigoplus_{p_i \mid |G_{n,1}|} \bigoplus_{j=1}^{r_i^{(n)}} \mathbb{Z}_{d_{i,j}^{(n)}}.$$

We define canonical generators $g_{i,j}^{(n)}$ of G_n by

$$\begin{aligned} g_{0,j}^{(n)} &= (0 \oplus \cdots \oplus 0 \oplus 1_j \oplus 0 \oplus \cdots \oplus 0) \oplus 0_{G_{n,1}}, \quad j = 1, 2, \dots, r_0^{(n)}, \\ g_{i,j}^{(n)} &= 0_{G_{n,0}} \oplus (0 \oplus \cdots \oplus 0 \oplus 1_{i,j} \oplus 0 \oplus \cdots \oplus 0), \quad j = 1, 2, \dots, r_i^{(n)}. \end{aligned}$$

For a finitely generated abelian group G_n , set

$$S^{(n)} = \{(0, 0), (0, 1)\} \cup \{(i, j); i \in \mathbb{N} \text{ with } p_i \mid |G_{n,1}|, j = 1, 2, \dots, r_i^{(n)}\}.$$

Let $I_{i,j}^{(n)}$ be a copy of the interval $[0, 2\pi]$ for each $(i, j) \in S^{(n)}$ and denote by $\iota_{i,j}^{(n)}$ the identity map from $[0, 2\pi]$ onto $I_{i,j}^{(n)}$, and let $T_j^{(n)}$ be a copy of the torus $\mathbb{T} = \{e^{it}; t \in [0, 2\pi]\}$ for each $j = 1, 2, \dots, r_0^{(n)}$ and denote by $\tau_j^{(n)}$ the identity map from $\mathbb{T}$ onto $T_j^{(n)}$. Set $z(t) = e^{it} \in \mathbb{T}$, $t \in \mathbb{R}$. We define a one-point union X_n by

$$X_n = I_{0,0}^{(n)} \vee I_{0,1}^{(n)} \vee \bigvee_{j=1}^{r_0^{(n)}} T_j^{(n)} \vee \bigvee_{p_i \mid |G_{n,1}|} \bigvee_{j=1}^{r_i^{(n)}} I_{i,j}^{(n)},$$

identifying the base point of X_n with $\iota_{i,j}^{(n)}(0)$, $(i, j) \in S^{(n)}$ and $\tau_j^{(n)}(z(0))$, $j = 1, 2, \dots, r_0^{(n)}$. Set $c_n = \tau_j^{(n)}(z(0)) = \iota_{i,j}^{(n)}(0) \in X_n$ and $a_{i,j}^{(n)} = \iota_{i,j}^{(n)}(2\pi) \in X_n$ for $(i, j) \in S^{(n)}$.

For a finitely generated abelian group G_n and relatively prime natural numbers $\tilde{p}_0$ and $\tilde{p}_1$ such that $d_{i,j}^{(n)} \mid \tilde{p}_0 \cdot \tilde{p}_1$ for any $(i, j) \in S^{(n)}$ with $i \geq 1$, we set $\natural(i) = 1 - i$ for $i = 0, 1$ and $\natural(\mu, \nu) = \natural(i)$ for $(\mu, \nu) \in S^{(n)}$ with $\mu \geq 1$ and $p_\mu \mid \tilde{p}_i$, and we define a C^* -algebra $A(G_n, \tilde{p}_0, \tilde{p}_1)$ by

$$\begin{aligned} A(G_n, \tilde{p}_0, \tilde{p}_1) &= \{f \in C(X_n) \otimes M_{\tilde{d}}; f(a_{0,i}^{(n)}) \in \epsilon_{0,i}^{(n)}(M_{\tilde{p}_0}), i = 0, 1, \\ &\quad f(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n)}(M_{\tilde{d}/d_{i,j}^{(n)}}), (i, j) \in S^{(n)} \text{ with } i \geq 1\}, \end{aligned}$$

where $\tilde{d} = \tilde{p}_0 \cdot \tilde{p}_1$ (and $d_{i,j}^{(n)} = p_i^{k_{i,j}^{(n)}}$) and $\epsilon_{0,i}^{(n)}$ and $\epsilon_{i,j}^{(n)}$ are unital embeddings from $M_{\tilde{p}_i}$ and $M_{\tilde{d}/d_{i,j}^{(n)}}$ onto $M_{\tilde{d}}$. It is not so hard to show that the C^* -algebra

$A(G_n, \tilde{p}_0, \tilde{p}_1)$ is independent of choices of $\epsilon_{i,j}^{(n)}$, $(i, j) \in S^{(n)}$. In this paper we say that $A(G_n, \tilde{p}_0, \tilde{p}_1)$ is a *dimension drop algebra* for G_n , $\tilde{p}_0$, and $\tilde{p}_1$. In particular we denote by $I(\tilde{p}_0, \tilde{p}_1)$ the dimension drop algebra for $\{0\}$, $\tilde{p}_0$, and $\tilde{p}_1$ with $(\tilde{p}_0, \tilde{p}_1) = 1$, (i.e.,

$$I(\tilde{p}_0, \tilde{p}_1) = \{f \in C(I_{0,0}^{(n)} \cup I_{0,1}^{(n)}) \otimes M_{\tilde{d}}; f(a_{0,0}^{(n)}) \in M_{\tilde{p}_0} \otimes 1_{\tilde{p}_1}, f(a_{0,1}^{(n)}) \in 1_{\tilde{p}_0} \otimes M_{\tilde{p}_1}\},$$

that was defined in [6].

Let $e_{0,0}$ be a minimal projection of $M_{\tilde{d}}$. We define $f_{0,j}^{(n)} \in A(G_n, \tilde{p}_0, \tilde{p}_1)$ for $j = 1, 2, \dots, r_0^{(n)}$ and $f_{i,j}^{(n)} \in A(G_n, \tilde{p}_0, \tilde{p}_1)$ for $(i, j) \in S^{(n)}$ with $i \geq 1$ by

$$f_{0,j}^{(n)}(x) = \begin{cases} \exp(ite_{0,0}), & x = \tau_j^{(n)}(z(t)) \in T_j^{(n)}, t \in [0, 2\pi], \\ 1_{\tilde{d}}, & \text{otherwise,} \end{cases}$$

$$f_{i,j}^{(n)}(x) = \begin{cases} \exp(ite_{0,0}), & x = \iota_{i,j}^{(n)}(t) \in I_{i,j}^{(n)}, \quad t \in [0, 2\pi], \\ 1_{\tilde{d}}, & \text{otherwise.} \end{cases}$$

Proposition 2.1. *Let G_n , $\tilde{p}_i$, $i = 0, 1$, and $A(G_n, \tilde{p}_0, \tilde{p}_1)$ be as above. Then it follows that $(K_0(A(G_n, \tilde{p}_0, \tilde{p}_1)), [1]) \cong (\mathbb{Z}, 1)$ and that*

$$K_1(A(G_n, \tilde{p}_0, \tilde{p}_1)) \cong G_n,$$

under the identification determined by

$$[f_{0,j}^{(n)}]_{K_1(A(G_n, \tilde{p}_0, \tilde{p}_1))} \cong g_{0,j}^{(n)}, \quad j = 1, 2, \dots, r_0^{(n)}, \quad [f_{i,j}^{(n)}]_{K_1(A(G_n, \tilde{p}_0, \tilde{p}_1))} \cong g_{i,j}^{(n)},$$

for $(i, j) \in S^{(n)}$ with $i \geq 1$.

Proof. Set $A_n = A(G_n, \tilde{p}_0, \tilde{p}_1)$ and $I = I(\tilde{p}_0, \tilde{p}_1)$, then we have that $K_1(I) \cong 0$ from $(\tilde{p}_0, \tilde{p}_1) = 1$. Set

$$\begin{aligned} B_{n,0} &= \bigoplus_{(i,j) \in S^{(n)}, i \geq 1} M_{\tilde{d}/d_{i,j}^{(n)}}, \\ X_{n,0} &= \bigcup_{j=1}^{r_0^{(n)}} \tau_j^{(n)}(z((0, 2\pi))) \cup \bigcup_{(i,j) \in S^{(n)}, i \geq 1} \iota_{i,j}^{(n)}((0, 2\pi)) \subset X_n, \\ C_{n,0} &= C_0(X_{n,0}) \otimes M_{\tilde{d}}. \end{aligned}$$

Then we have the exact sequence

$$0 \longrightarrow C_{n,0} \xrightarrow{\rho} A_n \xrightarrow{\sigma} I \oplus B_{n,0} \longrightarrow 0,$$

which induces the exact sequence on K -groups

$$K_0(I \oplus B_{n,0}) \xrightarrow{\delta_0} K_1((C_{n,0})^\sim) \longrightarrow K_1(A_n) \longrightarrow 0,$$

where δ_0 is the exponential map.

By $(\tilde{p}_0, \tilde{p}_1) = 1$, it follows that $(K_0(I), [1]_{K_0(I)}) \cong (\mathbb{Z}, 1)$. Let $e_{i,j}$ be a minimal projection of $M_{\tilde{d}/d_{i,j}^{(n)}} \subset B_{n,0}$. It follows that $K_0(B_{n,0}) \cong \bigoplus_{(i,j) \in S^{(n)}, i \geq 1} \mathbb{Z}$ under the identification defined by

$$[e_{i,j}]_{K_0(B_{n,0})} \cong 0 \oplus \dots \oplus 0 \oplus 1_{i,j} \oplus 0 \oplus \dots \oplus 0,$$

and it follows that $K_1((C_{n,0})^\sim) \cong \bigoplus_{j=1}^{r_0^{(n)}} \mathbb{Z} \oplus \bigoplus_{(i,j) \in S^{(n)}, i \geq 1} \mathbb{Z}$ by the identification defined by

$$[f_{0,j}^{(n)}]_{K_1((C_{n,0})^\sim)} \cong (0 \oplus \dots \oplus 0 \oplus 1_j \oplus 0 \oplus \dots \oplus 0) \oplus 0_{\bigoplus_{i,j} \mathbb{Z}},$$

$$[f_{i,j}^{(n)}]_{K_1((C_{n,0})^\sim)} \cong 0_{\bigoplus_j \mathbb{Z}} \oplus (0 \oplus \cdots \oplus 0 \oplus 1_{i,j} \oplus 0 \oplus \cdots \oplus 0).$$

Define $h \in (A_n)_{\text{sa}}$ by

$$h(x) = \begin{cases} (1 - \frac{t}{2\pi})1_{\tilde{d}}, & x = \iota_{i,j}^{(n)}(t) \in I_{i,j}^{(n)}, \ i \geq 1, \\ 1_{\tilde{d}}, & \text{otherwise.} \end{cases}$$

Because of the definition of δ_0 and $\sigma(h) = 1_I \oplus 0_{B_{n,0}}$, we have that

$$\delta_0([1_I \oplus 0_{B_{n,0}}]_{K_0(I \oplus B_{n,0})}) = -[\exp(2\pi i h)]_{K_1((C_{n,0})^\sim)} \cong 0_{\bigoplus_j \mathbb{Z}} \oplus (\tilde{d} \oplus \cdots \oplus \tilde{d}),$$

under the above identification. Define $h_{i,j}^{(n)} \in (A_n)_{\text{sa}}$ for $(i,j) \in S^{(n)}$ with $i \geq 1$ by

$$h_{i,j}^{(n)}(x) = \begin{cases} \frac{t}{2\pi} \epsilon_{i,j}^{(n)}(e_{i,j}), & x = \iota_{i,j}^{(n)}(t) \in I_{i,j}^{(n)}, \ t \in [0, 2\pi], \\ 0_{\tilde{d}}, & \text{otherwise.} \end{cases}$$

Since $\sigma(h_{i,j}^{(n)}) = 0_I \oplus e_{i,j}$, it follows that

$$\begin{aligned} \delta_0([0_I \oplus e_{i,j}]) &= -[\exp(2\pi i h_{i,j}^{(n)})]_{K_1((C_{n,0})^\sim)} = -d_{i,j}^{(n)}[f_{i,j}^{(n)}] \\ &= 0_{\bigoplus_j \mathbb{Z}} \oplus (0 \oplus \cdots \oplus 0 \oplus -d_{i,j}^{(n)} \oplus 0 \oplus \cdots \oplus 0). \end{aligned}$$

Therefore it follows that

$$\begin{aligned} K_1(A_n) &\cong K_1((C_{n,0})^\sim)/\delta_0(K_0(I_n \oplus B_{n,0})) \\ &\cong \bigoplus_{j=1}^{r_0^{(n)}} \mathbb{Z} \oplus \bigoplus_{(i,j) \in S^{(n)}, \ i \geq 1} \mathbb{Z}_{d_{i,j}^{(n)}}, \end{aligned}$$

and that $[f_{0,j}^{(n)}]_{K_1(A_n)}$ and $[f_{i,j}^{(n)}]_{K_1(A_n)}$ correspond to $g_{0,j}^{(n)}$ and $g_{i,j}^{(n)}$ because of the above identification $K_1((C_{n,0})^\sim) \cong \bigoplus_j \mathbb{Z} \oplus \bigoplus_{i,j} \mathbb{Z}$.

Since $K_0(C_{n,0}) \cong 0$, we have the exact sequence

$$0 \longrightarrow K_0(A_n) \xrightarrow{\sigma} K_0(I \oplus B_{n,0}) \xrightarrow{\delta_0} K_1((C_{n,0})^\sim).$$

Then we have that $(K_0(A_n), [1_{A_n}]) \cong (\ker(\delta_0), [\sigma(1_{A_n})])$. Because $[\sigma(1_{A_n})] = 1 \oplus (\tilde{d}/d_{i,j}^{(n)})_{i,j}$ is the generator of $\ker(\delta_0)$ we conclude that $(\ker(\delta_0), [\sigma(1_{A_n})]) \cong (\mathbb{Z}, 1)$. $\blacksquare$

The following definitions are necessary to show the UHF-embeddability in Proposition 5.2 of the projectionless C^* -algebra which will be obtained in Theorem 3.2. For a finitely generated abelian group G_n , we denote by $\tilde{I}_{0,j}^{(n)}$, $j = 1, 2, \dots, r_0^{(n)}$, $r_0^{(n)}$ copies of the interval $[0, \pi]$, by $\tilde{I}_{1,j}^{(n)}$, $r_0^{(n)}$ copies of the interval $[-\pi, 0]$, by $\tilde{\iota}_{0,j}^{(n)}$ the identity map $[0, \pi] \rightarrow \tilde{I}_{0,j}^{(n)}$, and by $\tilde{\iota}_{1,j}^{(n)}$ the identity map $[-\pi, 0] \rightarrow \tilde{I}_{1,j}^{(n)}$. We define a one-point union $\tilde{X}_n$ by

$$\tilde{X}_n = I_{0,0}^{(n)} \vee I_{0,1}^{(n)} \vee \bigvee_{j=1}^{r_0^{(n)}} (\tilde{I}_{0,j}^{(n)} \vee \tilde{I}_{1,j}^{(n)}) \vee \bigvee_{p_i \mid |G_{n,1}|} \bigvee_{j=1}^{r_i^{(n)}} I_{i,j}^{(n)},$$

identifying the base point of $\tilde{X}_n$ with $\iota_{i,j}^{(n)}(0)$ and $\tilde{\iota}_{i,j}^{(n)}(0)$. We set $b_{i,j}^{(n)} = \tilde{\iota}_{i,j}^{(n)}((-1)^i \pi)$ for $i = 0, 1$ and $j = 1, 2, \dots, r_0^{(n)}$, and we identify $A(G_n, \tilde{p}_0, \tilde{p}_1)$ with the C^* -subalgebra $\{f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}; f(b_{0,j}^{(n)}) = \epsilon_j^{(n)}(f(b_{1,j}^{(n)})), f(a_{0,i}^{(n)}) \in \epsilon_{0,i}^{(n)}(M_{\tilde{p}_i}), f(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n)}(M_{\tilde{d}/d_{i,j}^{(n)}})\}$, where $\epsilon_j^{(n)}$, $j = 1, 2, \dots, r_0^{(n)}$ are fixed automorphisms of $M_{\tilde{d}}$, and we denote the set of end points of $\tilde{X}_n$ by

$$E_n = \{a_{i,j}^{(n)}; (i, j) \in S^{(n)}\} \cup \{b_{i,j}^{(n)}; i = 0, 1, j = 1, 2, \dots, r_0^{(n)}\}.$$

Given G_n and $A(G_n, \tilde{p}_0, \tilde{p}_1)$ as in Proposition 2.1 let γ_n be an injective homomorphism from G_n into another finitely generated abelian group G_{n+1} . We shall show how to lift γ_n to an embedding of $A(G_n, \tilde{p}_0, \tilde{p}_1)$ into another dimension drop algebra $A(G_{n+1}, \tilde{p}'_0, \tilde{p}'_1)$.

Proposition 2.2. *Let γ_n be an injective group homomorphism from G_n into G_{n+1} . Then there exist relatively prime numbers $\tilde{p}'_0$ and $\tilde{p}'_1$ such that $d_{i',j'}^{(n+1)} | \tilde{p}'_0 \tilde{p}'_1 = \tilde{d}'$ for any $(i', j') \in S^{(n+1)}$ with $i' \geq 1$, automorphisms $\{\epsilon_{j'}^{(n+1)} \in \text{Aut}(M_{\tilde{d}'}); j = 1, 2, \dots, r_0^{(n+1)}\}$, unital embeddings $\{\epsilon_{i',j'}^{(n+1)} : M_{\tilde{d}'/d_{i',j'}^{(n+1)}} \hookrightarrow M_{\tilde{d}'}; (i', j') \in S^{(n+1)}\}$, where $d_{0,i}^{(n+1)} = \tilde{p}'_{\mathfrak{t}(i)}$ for $i = 0, 1$, and a unital embedding $\psi_n : C(\tilde{X}_n) \otimes M_{\tilde{d}} \hookrightarrow C(\tilde{X}_{n+1}) \otimes M_{\tilde{d}'}$ satisfying the following conditions:*

(0,0) $\psi_n(A(G_n, \tilde{p}_0, \tilde{p}_1)) \subset A(G_{n+1}, \tilde{p}'_0, \tilde{p}'_1)$ and $(\psi_n|_{A(G_n, \tilde{p}_0, \tilde{p}_1)})_* \cong \gamma_n$ under the identification that was defined in Proposition 2.1, where $A(G_{n+1}, \tilde{p}'_0, \tilde{p}'_1)$ are defined by $\epsilon_{j'}^{(n+1)}$ and $\epsilon_{i',j'}^{(n+1)}$, and $(\psi_n|_{A_n})_*$ is the induced map on K_1 -groups,

(0,1) for $\varepsilon > 0$ and $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$, if $\psi_n(f)^{-1}(\{0\})$ is ε -dense in $\tilde{X}_{n+1}$ then $f^{-1}(\{0\})$ is ε -dense in $\tilde{X}_n$,

(0,2) for any $\varepsilon > 0$ and $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$,

$$\begin{aligned} & \max\{|\text{Tr}_{\tilde{d}'}(\psi_n(f)(x) - \psi_n(f)(y))|; x, y \in \tilde{X}_{n+1}, d(x, y) \leq \varepsilon\} \\ & \leq \max\{|\text{Tr}_{\tilde{d}}(f(x) - f(y))|; x, y \in \tilde{X}_n, d(x, y) \leq \varepsilon\}, \end{aligned}$$

where d is the canonical metric on $\tilde{X}_n$,

(0,3) $\psi_n(\chi(E_n) \otimes 1_{\tilde{d}}) \leq \chi(E_{n+1}) \otimes 1_{\tilde{d}'}$.

Proof. Suppose that G_n and G_{n+1} are defined by $(r_i^{(n)}, d_{i,j}^{(n)})$ and $(r_i^{(n+1)}, d_{i,j}^{(n+1)})$ as in the beginning of this section and let $\gamma_n : G_n \rightarrow G_{n+1}$ be an injective group homomorphism. For the free part of G_n and $j = 1, 2, \dots, r_0^{(n)}$, there exist $\mu_{j',j} \in \mathbb{Z}$, $j' = 1, 2, \dots, r_0^{(n+1)}$ and $\nu_{i',j',0,j} \in \mathbb{Z}_+$, $(i', j') \in S^{(n+1)}$ with $i' \geq 1$ such that

$$\gamma_n(g_{0,j}^{(n)}) = \sum_{j'=1}^{r_0^{(n+1)}} \mu_{j',j} g_{0,j'}^{(n+1)} + \sum_{(i',j') \in S^{(n+1)}, i' \geq 1} \nu_{i',j',0,j} g_{i',j'}^{(n+1)},$$

and that $0 \leq \nu_{i',j',0,j} < d_{i',j'}^{(n+1)} (= p_{i',j'}^{k_{i',j'}^{(n+1)}})$. For the torsion group $G_{n,1}$ and $(i,j) \in S^{(n)}$ with $i \geq 1$, because of $\gamma_n(G_{n,p_i}) \subset G_{n+1,p_i}$, there exist $\nu_{j',i,j} \in \mathbb{Z}$, $j' = 1, 2, \dots, r_{i'}^{(n+1)}$ such that

$$\gamma_n(g_{i,j}^{(n)}) = \sum_{j'=1}^{r_i^{(n+1)}} \nu_{j',i,j} g_{i,j'}^{(n+1)},$$

and that $0 \leq \nu_{j',i,j} < d_{i,j'}^{(n+1)}$. When $p_{i'} \nmid |G_{n,1}|$ we define $r_{i'}^{(n)} = 0$, $d_{i',j}^{(n)} = 1$ and $\nu_{j',i',j} = 0$. Remark that $d_{i',j'}^{(n+1)} | d_{i',j}^{(n)} \nu_{j',i',j}$ (by $0 = \gamma_n(d_{i',j}^{(n)} g_{i',j}^{(n)}) = \sum d_{i',j}^{(n)} \nu_{j',i',j} g_{i',j'}^{(n+1)}$).

Let P_0 and P_1 be large natural numbers such that

$$P_0 P_1 > \max\left(\left\{\sum_{j=1}^{r_0^{(n)}} |\mu_{j'j}|; j' = 1, 2, \dots, r_0^{(n+1)}\right\} \cup \{(r_0^{(n)} + r_{i'}^{(n)}) d_{i',j'}^{(n+1)}; i', j'\},\right)$$

$$(P_0 \tilde{p}_0, P_1 \tilde{p}_1) = 1, \quad \text{and } d_{i',j'}^{(n+1)} | P_0 P_1,$$

for any $(i', j') \in S^{(n+1)}$ with $i' \geq 1$, and set $\tilde{p}'_i = P_i \tilde{p}_i$ for $i = 0, 1$, $\tilde{d}' = \tilde{p}'_0 \cdot \tilde{p}'_1$, and $m = P_0 P_1$. We define a unital embedding $\psi_n : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_{n+1}) \otimes M_{\tilde{d}'}$ by dividing into the following three cases depending on $x \in \tilde{X}_{n+1}$: For $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$ and for $x = \iota_{0,i}^{(n+1)}(t) \in I_{0,i}^{(n+1)}$, $i = 0, 1$, and $t \in [0, 2\pi]$ we define

$$\psi_n(f)(x) = f(\iota_{0,i}^{(n)}(t)) \otimes 1_m;$$

For $x = \tilde{\iota}_{i,j'}^{(n+1)}((-1)^i t) \in \tilde{I}_{i,j'}^{(n+1)}$, $i = 0, 1$, $j' = 1, 2, \dots, r_0^{(n)}$, and $t \in [0, \pi]$ we define

$$\psi_n(f)(x) = \bigoplus_{j=1}^{r_0^{(n)}} f(\tilde{\iota}_{\natural(i,j,j'),j}^{(n)}((-1)^{\natural(i,j,j')} t)) \otimes 1_{|\mu_{j',j}|} \oplus f(c_n) \otimes 1_{m_{0,j'}},$$

where $\natural(i, j, j') \in \{0, 1\}$ is defined by $\arg(\mu_{j',j})(-1)^i = (-1)^{\natural(i,j,j')}$ and where $m_{0,j'} = m - \sum_{j=1}^{r_0^{(n)}} |\mu_{j',j}|$; For $x = \iota_{i',j'}^{(n+1)}(t) \in I_{i',j'}^{(n+1)}$, $(i', j') \in S^{(n+1)}$ with $i' \geq 1$ and $t \in [0, 2\pi]$, we define

$$\begin{aligned} \psi_n(f)(x) &= \bigoplus_{j=1}^{r_0^{(n)}} (f(\tilde{\iota}_{0,j}^{(n)}(t/2)) \otimes 1_{\nu_{i',j',0,j}} \oplus f(\tilde{\iota}_{1,j}^{(n)}(-t/2)) \otimes 1_{d_{i',j'}^{(n+1)} - \nu_{i',j',0,j}}) \\ &\oplus \bigoplus_{j=1}^{r_{i'}^{(n)}} (f(\iota_{i',j}^{(n)}(t)) \otimes 1_{\nu_{j',i',j}} \oplus f(\iota_{0,\natural(i',j,j')}^{(n)}(t)) \otimes 1_{d_{i',j'}^{(n+1)} - \nu_{j',i',j}}) \\ &\oplus f(c_n) \otimes 1_{m - (r_0^{(n)} + r_{i'}^{(n)}) d_{i',j'}^{(n+1)}}. \end{aligned}$$

Note that when $p_{i'} \nmid |G_{n,1}|$ we have defined $r_{i'}^{(n)} = 0$. It is not so hard to show that $\psi_n(f)(x) \rightarrow f(c_n) \otimes 1_m$, $x \rightarrow c_{n+1}$ for $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$. Thus it follows that $\psi_n(C(\tilde{X}_n) \otimes M_{\tilde{d}}) \subset C(\tilde{X}_{n+1}) \otimes M_{\tilde{d}'}$.

Recall that $A_n = A(G_n, \tilde{p}_0, \tilde{p}_1)$ was defined in terms of automorphisms $\epsilon_j^{(n)}$ and also depend on unital embeddings $\epsilon_{i,j}^{(n)}$. We shall show that there exist automorphisms $\epsilon_{j'}^{(n+1)}$ of $M_{\tilde{d}'}$ and unital embeddings $\epsilon_{i',j'}^{(n+1)}$ which satisfy the condition in (0,0). Let $\epsilon_{j'}^{(n+1)}$ be an extension of $(\bigoplus_{j=1}^{r_0^{(n)}} (\epsilon_j^{(n)})^{\arg(\mu_{j',j})} \otimes 1_{|\mu_{j',j}|}) \oplus \text{id}_{M_{\tilde{d}}} \otimes 1_{m_{0,j'}}$, then we have $\epsilon_{j'}^{(n+1)}(\psi_n(f)(b_{1,j'}^{(n+1)})) = \psi_n(f)(b_{0,j'}^{(n+1)})$ for $j' = 1, 2, \dots, r_0^{(n+1)}$. Since $\tilde{p}_{\mathfrak{h}(i)} | m\tilde{p}_{\mathfrak{h}(i)}$, $i = 0, 1$, $f(a_{0,i}^{(n)}) \in \epsilon_{0,i}^{(n)}(M_{\tilde{p}_i})$ for any $f \in A_n$, and $\psi_n(f)(a_{0,i}^{(n+1)}) = f(a_{0,i}^{(n)}) \otimes 1_m$ there exists a unital embedding $\epsilon_{0,i}^{(n+1)} : M_{\tilde{p}_i} \hookrightarrow M_{\tilde{d}'}$ such that

$$\psi_n(f)(a_{0,i}^{(n+1)}) \in \epsilon_{0,i}^{(n+1)}(M_{\tilde{p}_i}), \quad f \in A_n, \quad i = 0, 1.$$

By the definition of ψ_n , it follows that for $f \in A_n$,

$$\begin{aligned} \psi_n(f)(a_{i',j'}^{(n+1)}) &= \bigoplus_{j=1}^{r_0^{(n)}} \epsilon_j^{(n)}(f(b_{1,j}^{(n)})) \otimes 1_{\nu_{i',j',0,j}} \oplus f(b_{1,j}^{(n)}) \otimes 1_{d_{i',j'}^{(n+1)} - \nu_{i',j',0,j}} \\ &\oplus \bigoplus_{j=1}^{r_{i'}^{(n)}} f(a_{i',j}^{(n)}) \otimes 1_{\nu_{j',i',j}} \oplus f(a_{0,\mathfrak{h}(i',j')}^{(n)}) \otimes 1_{d_{i',j'}^{(n+1)} - \nu_{j',i',j}} \\ &\oplus f(c_n) \otimes 1_{m - (r_0^{(n)} + r_{i'}^{(n)})d_{i',j'}^{(n+1)}}, \end{aligned}$$

Since $d_{i',j'}^{(n+1)} | d_{i',j}^{(n)} \nu_{j',i',j}$ for any $j = 1, 2, \dots, r_{i'}^{(n)}$ and since $f(a_{i',j}^{(n)}) \in \epsilon_{i',j}^{(n)}(M_{\tilde{d}/d_{i',j}^{(n)}})$ for $f \in A_n$, we obtain a unitary $u_{i',j',j}$ in $M_{\tilde{d}\nu_{j',i',j}}$ such that

$$\text{Ad } u_{i',j',j}(f(a_{i',j}^{(n)}) \otimes 1_{\nu_{j',i',j}}) \in M_{\tilde{d}\nu_{j',i',j}/d_{i',j'}^{(n+1)}} \otimes 1_{d_{i',j'}^{(n+1)}}, \quad f \in A_n, \quad p_{i'} | |G_{n,1}|.$$

When $p_{i'} \nmid |G_{n,1}|$ we have defined $\nu_{j',i',j} = 0$. When $p_{i'} | |G_{n,1}|$ we have $d_{i',j}^{(n)} | \tilde{p}_{\mathfrak{h}(i',j)}$, then it follows that $d_{i',j'}^{(n+1)} | \tilde{p}_{\mathfrak{h}(i',j)}(d_{i',j'}^{(n+1)} - \nu_{j',i',j})$, and from $f(a_{0,\mathfrak{h}(i',j)}^{(n)}) \in \epsilon_{0,\mathfrak{h}(i',j)}^{(n)}(M_{\tilde{p}_{\mathfrak{h}(i',j)}})$ for $f \in A_n$ we obtain a unitary $v_{i',j',j}$ in $M_{\tilde{d}(d_{i',j'}^{(n+1)} - \nu_{j',i',j})}$ such that

$$\text{Ad } v_{i',j',j}(f(a_{0,\mathfrak{h}(i',j)}^{(n)}) \otimes 1_{d_{i',j'}^{(n+1)} - \nu_{j',i',j}}) \in M_N \otimes 1_{d_{i',j'}^{(n+1)}}, \quad f \in A_n,$$

where $N = \tilde{d}(d_{i',j'}^{(n+1)} - \nu_{j',i',j})/d_{i',j'}^{(n+1)}$. Hence there exists a unital embedding $\epsilon_{i',j'}^{(n+1)} : M_{\tilde{d}'/d_{i',j'}^{(n+1)}} \hookrightarrow M_{\tilde{d}'}$ such that

$$\psi_n(f)(a_{i',j'}^{(n+1)}) \in \epsilon_{i',j'}^{(n+1)}(M_{\tilde{d}'/d_{i',j'}^{(n+1)}}), \quad f \in A_n.$$

Whence we conclude that $\psi_n(A_n) \subset A(G_{n+1}, \tilde{p}_0, \tilde{p}_1)$.

Proof of $(\psi_n|_{A_n})_* \cong \gamma_n$. In Proposition 2.1, we have identified $[f_{0,j}^{(n)}]_{K_1(A_n)}$ and $[f_{i,j}^{(n)}]_{K_1(A_n)}$ with $g_{0,j}^{(n)}$ and $g_{i,j}^{(n)}$. Set $A_{n+1} = A(G_{n+1}, \tilde{p}_0, \tilde{p}_1)$ and

$$f'_{0,j}(x) = \begin{cases} \exp(2ite_{0,0}), & x = \tilde{t}_{0,j}^{(n)}(t) \in \tilde{I}_{0,j}^{(n)}, \quad 0 \leq t \leq \pi, \\ 1_{\tilde{d}}, & \text{otherwise,} \end{cases}$$

where $e_{0,0}$ is a minimal projection of $M_{\tilde{d}}$. Remark that $f'_{0,j} \in U(A_n)$ and $[f'_{0,j}]_{K_1(A_n)} = [f_{0,j}^{(n)}] = g_{0,j}^{(n)}$. By the definition of ψ_n it follows that $\psi_n(f'_{0,j})(x)$

$$= \begin{cases} \exp(2ite_{0,0} \otimes 1_{|\mu_{j',j}|}), & x = \tilde{t}_{(0,j,j'),j'}^{(n+1)}(\arg(\mu_{j',j})t), \quad 0 \leq t \leq \pi, \\ \exp(ite_{0,0} \otimes 1_{\nu_{i',j',0,j}}), & x = \tilde{t}_{i',j'}^{(n+1)}(t) \in I_{i',j'}^{(n+1)}, \quad i' \geq 1, \quad 0 \leq t \leq 2\pi, \\ 1_{\tilde{d}}, & \text{otherwise,} \end{cases}$$

which implies that $\psi_n(f'_{0,j}) \in U(A_{n+1})$ and

$$[\psi_n(f'_{0,j})]_{K_1(A_{n+1})} = \sum_{j'} \mu_{j',j} g_{0,j'}^{(n+1)} + \sum_{i'} \sum_{j'} \nu_{i',j',0,j} g_{i',j'}^{(n+1)}.$$

For $(i,j) \in S^{(n)}$ with $i \geq 1$ it follows that

$$\psi_n(f_{i,j}^{(n)})(x) = \begin{cases} \exp(ite_{0,0} \otimes 1_{\nu_{j',i,j}}), & x = \tilde{t}_{i,j'}^{(n+1)}(t) \in I_{i,j'}^{(n+1)}, \\ 1_{\tilde{d}}, & \text{otherwise,} \end{cases}$$

which implies that $\psi_n(f_{i,j}^{(n)}) \in U(A_{n+1})$ and

$$[\psi_n(f_{i,j}^{(n)})]_{K_1(A_{n+1})} = \sum_{j'} \nu_{j',i,j} g_{i,j'}^{(n+1)}.$$

Hence we have that $(\psi_n)_* \cong \gamma_n$.

Proof of $(0,1)$. Let $\hat{\psi}_{n,\delta} : \tilde{X}_{n+1} \rightarrow \tilde{X}_n$, $\delta = 1, 2, \dots, m$ be the continuous maps

such that $\bigoplus_{\delta=1}^m f \circ \hat{\psi}_{n,\delta}(x) = \psi_n(f)(x)$, $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$. Since γ_n is injective, for

$j_0 \in \{1, 2, \dots, r_0^{(n)}\}$, there exists $j'_0 \in \{1, 2, \dots, r_0^{(n+1)}\}$ such that $\mu_{j'_0, j_0} \neq 0$. Set

$\delta_0 = \sum_{j=1}^{j_0-1} |\mu_{j'_0, j}| + 1$. By the definition of ψ_n at $x = \tilde{t}_{i,j'_0}((-1)^i t)$, we have that

$\hat{\psi}_{n,\delta_0}(\tilde{I}_{0,j'_0}^{(n+1)} \cup \tilde{I}_{1,j'_0}^{(n+1)}) = \tilde{I}_{0,j_0}^{(n)} \cup \tilde{I}_{1,j_0}^{(n)}$ and that if $\psi_n(f)^{-1}(\{0\}) \cap (\tilde{I}_{0,j'_0}^{(n+1)} \cup \tilde{I}_{1,j'_0}^{(n+1)})$ is ε -dense then $\hat{\psi}_{n,\delta_0}(\psi_n(f)^{-1}(\{0\}) \cap (\tilde{I}_{0,j'_0}^{(n+1)} \cup \tilde{I}_{1,j'_0}^{(n+1)}))$ is ε -dense in $\tilde{I}_{0,j_0}^{(n)} \cup \tilde{I}_{1,j_0}^{(n)}$. Since $f^{-1}(\{0\}) \cap (\tilde{I}_{0,j_0}^{(n)} \cup \tilde{I}_{1,j_0}^{(n)}) \supset \hat{\psi}_{n,\delta_0}(\psi_n(f)^{-1}(\{0\}) \cap (\tilde{I}_{0,j'_0}^{(n+1)} \cup \tilde{I}_{1,j'_0}^{(n+1)}))$, it follows that if $\psi_n(f)^{-1}(\{0\}) \cap (\tilde{I}_{0,j'_0}^{(n+1)} \cup \tilde{I}_{1,j'_0}^{(n+1)})$ is ε -dense then $f^{-1}(\{0\}) \cap (\tilde{I}_{0,j_0}^{(n)} \cup \tilde{I}_{1,j_0}^{(n)})$ is ε -dense. Similarly, since γ_n is injective, for $(i_0, j_0) \in S^{(n)}$ there exists

$(i_0, j'_0) \in S^{(n+1)}$ such that $\nu_{j'_0, i_0, j_0} \neq 0$. Set $\delta_0 = (r_0^{(n)} + j_0 - 1)d_{i_0, j'_0}^{(n+1)} + 1$. By the definition of ψ_n at $x = \iota_{i_0, j'_0}^{(n+1)}(t)$, we have that $\hat{\psi}_{n, \delta_0}(I_{i_0, j'_0}^{(n+1)}) = I_{i_0, j_0}^{(n)}$ and that if $\psi_n(f)^{-1}(\{0\}) \cap I_{i_0, j'_0}^{(n+1)}$ is ε -dense then $\hat{\psi}_{n, \delta_0}(\psi_n(f)^{-1}(\{0\}) \cap I_{i_0, j'_0}^{(n+1)}) \subset f^{-1}(\{0\}) \cap I_{i_0, j_0}^{(n)}$ is ε -dense. Considering a neighborhood of c_n , we notice that if $\psi_n(f)^{-1}(\{0\}) \cap (\tilde{X}_{n+1} \setminus (I_{0,0}^{(n+1)} \cup I_{0,1}^{(n+1)})) \neq \emptyset$ then $f(c_n) = 0$. If $\psi_n(f)^{-1}(\{0\}) \cap (\tilde{X}_{n+1} \setminus (I_{0,0}^{(n+1)} \cup I_{0,1}^{(n+1)})) = \emptyset$ and $\psi_n(f)^{-1}(\{0\})$ is ε -dense then there exists $(0, j) \in S^{(n+1)}$ such that $d(\psi_n(f)^{-1}(\{0\}) \cap I_{0,j}^{(n+1)}, c_{n+1}) \leq \varepsilon$, which induces that $d(f^{-1}(\{0\}) \cap I_{0,j}^{(n)}, c_n) \leq \varepsilon$, $j = 0$ or 1 . Hence, combining the above arguments, we have (0,1).

Proof of (0,2). Let $\hat{\psi}_{n, \delta} : \tilde{X}_{n+1} \rightarrow \tilde{X}_n$, $\delta = 1, 2, \dots, m$ be the continuous maps such that $\bigoplus_{\delta=1}^m f \circ \hat{\psi}_{n, \delta}(x) = \psi_n(f)(x)$, $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$. By the definition of ψ_n , we have that

$$d(\hat{\psi}_{n, \delta}(x), \hat{\psi}_{n, \delta}(y)) \leq d(x, y),$$

for any $x, y \in \tilde{X}_{n+1}$ and $\delta = 1, 2, \dots, m$. Then for x and $y \in \tilde{X}_{n+1}$ with $d(x, y) \leq \varepsilon$ and $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$, it follows that

$$\begin{aligned} & |\text{Tr}_{\tilde{d}'}(\psi_n(f)(x) - \psi_n(f)(y))| \\ & \leq \frac{1}{m} \sum_{\delta=1}^m |\text{Tr}_{\tilde{d}}(f(\hat{\psi}_{n, \delta}(x)) - f(\hat{\psi}_{n, \delta}(y)))| \\ & \leq \max\{|\text{Tr}(f(x) - f(y))|; x, y \in \tilde{X}_n, d(x, y) \leq \varepsilon\}. \end{aligned}$$

Thus we have (0,2).

Proof of (0,3). By the definition of ψ_n , for $h_k^{(n)} \in C(\tilde{X}_n)_+^1$, $n, k \in \mathbb{N}$ with $h_k^{(n)} \searrow \chi(E_n)$ ($k \rightarrow \infty$), ψ_n satisfies that

$$\lim_{k \rightarrow \infty} \psi_n(h_k^{(n)} \otimes 1_{\tilde{d}}) \leq \chi(E_{n+1}) \otimes 1_{\tilde{d}}.$$

Then we have (0,3). This completes the proof of the proposition. $\blacksquare$

3 Simple projectionless C^* -algebras with a unique tracial state

In this section, we construct the inductive limit C^* -algebra of a sequence of dimension drop algebras that is simple and has a unique tracial state.

For the simplicity of C^* -algebras and the uniqueness of tracial states, we prepare the following proposition.

Proposition 3.1. *Let G_n be a finitely generated abelian group, $\tilde{p}_0$ and $\tilde{p}_1$ relatively prime numbers such that $d_{i,j}^{(n)} | \tilde{d} = \tilde{p}_0 \cdot \tilde{p}_1$ for any $(i, j) \in S^{(n)}$ with*

$i \geq 1$, A_n a dimension drop algebra for G_n , $\tilde{p}_0$, and $\tilde{p}_1$. Then there exist relatively prime numbers $\tilde{p}'_0$ and $\tilde{p}'_1$ such that $\tilde{p}_0 | \tilde{p}'_0$ and $\tilde{p}_1 | \tilde{p}'_1$, automorphisms $\{\epsilon_j^{(n,0)} \in \text{Aut}(M_{\tilde{d}'}); j = 1, 2, \dots, r_0^{(n)}\}$, where $\tilde{d}' = \tilde{p}'_0 \tilde{p}'_1$, unital embeddings $\{\epsilon_{i,j}^{(n,0)} : M_{\tilde{d}'/d_{i,j}^{(n)}} \hookrightarrow M_{\tilde{d}'}; (i,j) \in S^{(n)}\}$, where $d_{0,i}^{(n)} = \tilde{p}'_{\mathfrak{h}(i)}$ for $i = 0, 1$, and a unital embedding $\varphi_{n,0} : C(\tilde{X}_n) \otimes M_{\tilde{d}} \hookrightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}'}$ satisfying the following conditions:

- (0) $\varphi_{n,0}(A_n) \subset A(G_n, \tilde{p}'_0, \tilde{p}'_1)$ and $(\varphi_{n,0}|_{A_n})_* \cong \text{id}_{G_n}$ under the identification which is defined in Proposition 2.1, where $A(G_n, \tilde{p}'_0, \tilde{p}'_1)$ is defined by $\epsilon_j^{(n,0)}$ and $\epsilon_{i,j}^{(n,0)}$,
- (1) if $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$ satisfies that $\varphi_{n,0}(f)^{-1}(\{0\})$ is ε -dense in $\tilde{X}_n$ then $f^{-1}(\{0\})$ is $1/2\varepsilon$ -dense in $\tilde{X}_n$,
- (2) for any $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$ and $\varepsilon > 0$,

$$\begin{aligned} & \max\{|\text{Tr}_{\tilde{d}'}(\varphi_{n,0}(f)(x) - \varphi_{n,0}(f)(y))|; x, y \in \tilde{X}_n, d(x, y) \leq \varepsilon\} \\ & \leq \max\{|\text{Tr}_{\tilde{d}}(f(x) - f(y))|; x, y \in \tilde{X}_n, d(x, y) \leq 1/2\varepsilon\}, \end{aligned}$$

where d is the canonical metric on $\tilde{X}_n$,

- (3) $\text{Tr}_{\tilde{d}'}(\varphi_{n,0}(\chi(E_n) \otimes 1_{\tilde{d}})(x)) \leq 1/2 \text{Tr}(\chi(E_n) \otimes 1_{\tilde{d}'}(x)), \quad x \in \tilde{X}_n$.

Proof. Let G_n , $\tilde{p}_0$, $\tilde{p}_1$, and $\tilde{d}$ be as in Section 2 (i.e., by $r_0^{(n)} \in \mathbb{Z}_+$, $r_i^{(n)}, k_{i,j}^{(n)} \in \mathbb{N}$ for $i \geq 1$ with $p_i | |G_{n,1}|$, and $d_{i,j}^{(n)} = p_i^{k_{i,j}^{(n)}}$, we have represented G_n as

$$G_n \cong \mathbb{Z}^{r_0^{(n)}} \oplus \bigoplus_{p_i | |G_{n,1}|} \bigoplus_{j=1}^{r_i^{(n)}} \mathbb{Z}_{d_{i,j}^{(n)}},$$

($\tilde{p}_0, \tilde{p}_1 = 1$, $\tilde{d} = \tilde{p}_0 \cdot \tilde{p}_1$, and $d_{i,j}^{(n)} | \tilde{d}$ for any $(i, j) \in S^{(n)}$ with $i \geq 1$), and let A_n be a dimension drop algebra for G_n , $\tilde{p}_0$, and $\tilde{p}_1$.

First, we define $\tilde{p}'_i$, $i = 0, 1$ as follows. Let $P_i \in \mathbb{N}$, $i = 0, 1$ be such that $(P_0 \tilde{p}_0, P_1 \tilde{p}_1) = 1$, $P_i > \tilde{d}$, and

$$P_0 \equiv P_1 \equiv 1 \pmod{\tilde{d}}$$

($P_0 = a'\tilde{d} + 1$, $P_1 = b'P_0\tilde{d} + 1$, $a', b' \in \mathbb{N}$). We define $\tilde{p}'_i = P_i \tilde{p}_i$ for $i = 0, 1$, and $m = P_0 P_1$. By the definition of P_0 and P_1 , we obtain a, b and $c \in \mathbb{N}$ satisfying that

$$m = a\tilde{p}'_0 + P_0 = b\tilde{p}'_1 + P_1 = c\tilde{d} + 1.$$

Set

$$\kappa = P_0 + P_1 + \sum_{p_i | |G_{n,1}|} r_i^{(n)},$$

then, taking large P_i , we may assume that $m \geq 2(3r_0^{(n)} + \kappa)$.

We shall define $\varphi_{n,0}$ as the direct sum of $r_0^{(n)}$ homomorphisms of $C(\tilde{X}_n) \otimes M_{\tilde{d}}$ into $C(\tilde{X}_n) \otimes M_{\tilde{d}} \otimes M_3$, κ homomorphisms of $C(\tilde{X}_n) \otimes M_{\tilde{d}}$ into $C(\tilde{X}_n) \otimes M_{\tilde{d}}$, and the homomorphism ξ_c of $C(\tilde{X}_n) \otimes M_{\tilde{d}}$ into $C(\tilde{X}_n) \otimes M_{\tilde{d}} \otimes M_{m-(3r_0^{(n)}+\kappa)}$ defined by $f \mapsto f(c_n) \otimes 1_{m-(3r_0^{(n)}+\kappa)}$. The first type of homomorphisms, denoted by η_j , will be constructed to preserve the free part of G_n and the second type denoted by ξ_δ will be constructed to preserve the torsion subgroup of G_n . We define a *-homomorphism $\eta : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{3r_0^{(n)}\tilde{d}}$ as follows. Set

$$\begin{aligned} D(t) &= \text{diag}(z(\frac{t}{3}), z(\frac{t+2\pi}{3}), z(\frac{t-2\pi}{3})) \in U(M_3), \quad t \in [-\pi, \pi], \\ F(t) &= \text{diag}(0, \frac{2\pi-t}{3}, \frac{t-2\pi}{3}) \in M_3, \quad t \in [0, 2\pi]. \end{aligned}$$

and let $V \in U(M_3)$ be such that

$$VD(-\pi)V^* = D(\pi).$$

For $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$ and $j \in \{1, 2, \dots, r_0^{(n)}\}$ let $f \circ \tau_j^{(n)}(D(t))$

$$= \begin{cases} \text{diag}(f \circ \tilde{\iota}_{0,j}(\frac{t}{3}), f \circ \tilde{\iota}_{0,j}(\frac{t+2\pi}{3}), f \circ \tilde{\iota}_{1,j}(\frac{t-2\pi}{3})), & t \in [0, \pi] \\ \text{diag}(f \circ \tilde{\iota}_{1,j}(\frac{t}{3}), f \circ \tilde{\iota}_{0,j}(\frac{t+2\pi}{3}), f \circ \tilde{\iota}_{1,j}(\frac{t-2\pi}{3})), & t \in [-\pi, 0], \end{cases}$$

and let $f \circ \tau_j^{(n)} \circ z(F(t))$

$$= \text{diag}(f(c_n), f \circ \tilde{\iota}_{0,j}(\frac{2\pi-t}{3}), f \circ \tilde{\iota}_{1,j}(\frac{t-2\pi}{3})), \quad t \in [0, 2\pi].$$

Remark that for $f \in A_n$,

$$\text{Ad } 1_{\tilde{d}} \otimes V \cdot (1_{\tilde{d}} \oplus 1_{\tilde{d}} \oplus v_j^{(n)})(f \circ \tau_j^{(n)}(D(-\pi))) = f \circ \tau_j^{(n)}(D(\pi)),$$

where $v_j^{(n)}$ is a unitary in $M_{\tilde{d}}$ satisfying that $\text{Ad } v_j^{(n)}(f(b_{1,j}^{(n)})) = f(b_{0,j}^{(n)})$ for any $f \in A_n$. Set $v'_j = 1_{\tilde{d}} \otimes V \cdot (1 \oplus 1 \oplus v_j^{(n)})$.

Define *-homomorphisms $\eta_j : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}} \otimes M_3$, $j = 1, 2, \dots, r_0^{(n)}$ by

$$\eta_j(f)(x) = \begin{cases} f \circ \tau_j^{(n)}(D((-1)^i t)), & x = \tilde{\iota}_{i,j}^{(n)}((-1)^i t), \quad t \in [0, \pi], \quad i = 0, 1, \\ f \circ \tau_j^{(n)}(D(0)), & x \in \tilde{I}_{0,\nu}^{(n)} \vee \tilde{I}_{1,\nu}^{(n)}, \quad \nu \neq j, \\ f \circ \tau_j^{(n)} \circ z(F(t)), & x = \iota_{\mu,\nu}^{(n)}(t) \in I_{\mu,\nu}^{(n)}, \quad (\mu, \nu) \in S^{(n)}, \end{cases}$$

for $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$. Remark that

$$\eta_j(f)(a_{\mu,\nu}^{(n)}) = f(c_n) \otimes 1_3, \quad f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}, \quad (\mu, \nu) \in S^{(n)},$$

and that

$$\text{Ad } v'_j(\eta_j(f)(b_{1,j}^{(n)})) = \eta_j(f)(b_{0,j}^{(n)}), \quad f \in A_n, \quad j = 1, 2, \dots, r_0^{(n)}.$$

We define a $*$ -homomorphism $\eta : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{3r_0^{(n)}\tilde{d}}$ by $\eta = \bigoplus_{j=1}^{r_0^{(n)}} \eta_j$.

We define a $*$ -homomorphism $\xi_\delta^{(n)} : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}}$, $\delta = 1, 2, \dots, \kappa$ by the following cases : when $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$ and $1 \leq \delta \leq P_1$

$$\xi_\delta^{(n)}(f)(x) = \begin{cases} f(\iota_{0,0}^{(n)}(\pi + \frac{t}{2})), & x = \iota_{0,0}^{(n)}(t) \in I_{0,0}^{(n)}, \\ f(\iota_{0,0}^{(n)}(\pi - \frac{t}{2})), & x = \iota_{\mu,\nu}^{(n)}(t) \in I_{\mu,\nu}^{(n)}, \quad (\mu, \nu) \neq (0, 0), \\ f(\iota_{0,0}^{(n)}(\pi)), & x \in \tilde{I}_{0,j}^{(n)} \cup \tilde{I}_{1,j}^{(n)}, \quad j = 1, 2, \dots, r_0^{(n)}, \end{cases}$$

when $P_1 + 1 \leq \delta \leq P_0 + P_1$

$$\xi_\delta^{(n)}(f)(x) = \begin{cases} f(\iota_{0,1}^{(n)}(\pi + \frac{t}{2})), & x = \iota_{0,1}^{(n)}(t) \in I_{0,1}^{(n)}, \\ f(\iota_{0,1}^{(n)}(\pi - \frac{t}{2})), & x = \iota_{\mu,\nu}^{(n)}(t) \in I_{\mu,\nu}^{(n)}, \quad (\mu, \nu) \neq (0, 1), \\ f(\iota_{0,1}^{(n)}(\pi)), & x \in \tilde{I}_{0,j}^{(n)} \cup \tilde{I}_{1,j}^{(n)}, \quad j = 1, 2, \dots, r_0^{(n)}, \end{cases}$$

when $\delta = P_0 + P_1 + \sum_{\mu < i, \quad p_\mu \mid |G_{n,1}|} r_\mu^{(n)} + j, \quad p_i \mid |G_{n,1}|$, and $j = 1, 2, \dots, r_i^{(n)}$

$$\xi_\delta^{(n)}(f)(x) = \begin{cases} f(\iota_{i,j}^{(n)}(\pi + \frac{t}{2})), & x = \iota_{i,j}^{(n)}(t) \in I_{i,j}^{(n)}, \\ f(\iota_{i,j}^{(n)}(\pi - \frac{t}{2})), & x = \iota_{\mu,\nu}^{(n)}(t) \in I_{\mu,\nu}^{(n)}, \quad (\mu, \nu) \neq (i, j), \\ f(\iota_{i,j}^{(n)}(\pi)), & x \in \tilde{I}_{0,j}^{(n)} \cup \tilde{I}_{1,j}^{(n)}, \quad j = 1, 2, \dots, r_0^{(n)}, \end{cases}$$

We define a $*$ -homomorphism $\xi : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{\kappa\tilde{d}}$ by $\xi = \bigoplus_{\delta=1}^{\kappa} \xi_\delta$.

Set $\tilde{d}' = \tilde{p}_0' \tilde{p}_1' (= m\tilde{d})$, and let A_n be defined by automorphisms $\epsilon_j^{(n)} (= \text{Ad } v_j^{(n)})$ and unital embeddings $\epsilon_{i,j}^{(n)}$. Define a $*$ -homomorphism $\xi_c : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{(m-(3r_0^{(n)}+\kappa))\tilde{d}}$ by $\xi_c(f)(x) = f(c_n) \otimes 1_{m-(3r_0^{(n)}+\kappa)}$, $x \in \tilde{X}_n$, and define a unital $*$ -homomorphisms $\varphi_{n,0} : C(\tilde{X}_n) \otimes M_{\tilde{d}} \hookrightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}'}$ by

$$\varphi_{n,0} = \eta \oplus \xi \oplus \xi_c.$$

We shall show that there exist automorphisms $\epsilon_j^{(n,0)}$ and unital embeddings $\epsilon_{i,j}^{(n,0)}$ satisfying the condition in (0). Set $v_j = 1_{(j-1)\tilde{d}} \oplus v'_j \oplus 1_{(m-j)\tilde{d}} \in U(M_{\tilde{d}'})$ and $\epsilon_j^{(n,0)} = \text{Ad } v_j$, then it follows that

$$\epsilon_j^{(n,0)}(\varphi_{n,0}(f)(b_{1,j}^{(n)})) = \varphi_{n,0}(f)(b_{0,j}^{(n)}), \quad f \in A_n, \quad j = 1, 2, \dots, r_0^{(n)}.$$

From the remark of η and the definition of ξ it follows that

$$\varphi_{n,0}(f)(a_{0,i}^{(n)}) = f(c_n) \oplus \dots \oplus f(c_n) \oplus f(a_{0,i}^{(n)}) \otimes 1_{P_{\mathfrak{a}(i)}} \oplus f(c_n) \oplus \dots \oplus f(c_n),$$

for $i = 0, 1$. Since $f(a_{0,i}^{(n)}) \in \epsilon_{0,i}^{(n)}(M_{\tilde{p}_i})$ for $f \in A_n$, $\tilde{p}_{\natural(i)}' = P_{\natural(i)}\tilde{p}_{\natural(i)}$, and $\tilde{p}_{\natural(i)}'|m - P_{\natural(i)}$ there exist unital embeddings $\epsilon_{0,i}^{(n,0)} : M_{\tilde{p}_i'} \hookrightarrow M_{\tilde{d}'}$, $i = 0, 1$, such that

$$\varphi_{n,0}(f)(a_{0,i}^{(n)}) \in \epsilon_{0,i}^{(n,0)}(M_{\tilde{p}_i'}), \text{ for any } f \in A_n.$$

Similarly, it follows that

$$\varphi_{n,0}(f)(a_{i,j}^{(n)}) = f(c_n) \oplus \cdots \oplus f(c_n) \oplus f(a_{i,j}^{(n)}) \oplus f(c_n) \oplus \cdots \oplus f(c_n),$$

for any $(i, j) \in S^{(n)}$ with $i \geq 1$. Since $f(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n)}(M_{\tilde{d}/d_{i,j}^{(n)}})$ and $d_{i,j}^{(n)} | \tilde{d} | m - 1$, there exist unital embeddings $\epsilon_{i,j}^{(n,0)} : M_{\tilde{d}'/d_{i,j}^{(n)}} \hookrightarrow M_{\tilde{d}'}$ such that

$$\varphi_{n,0}(f)(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n,0)}(M_{\tilde{d}'/d_{i,j}^{(n)}}), \text{ for any } f \in A_n.$$

Hence we conclude that $\varphi_{n,0}(A_n) \subset A(G_n, \tilde{p}_0', \tilde{p}_1')$.

Proof of $(\varphi_{n,0}|_{A_n})_* = \text{id}_{G_n}$. Define $f'_{0,j} \in U(A_n)$ for $j = 1, 2, \dots, r_0^{(n)}$ by

$$f'_{0,j}(x) = \begin{cases} \exp(6ite_{0,0}), & x = \tilde{t}_{0,j}^{(n)}(t), \quad 0 \leq t \leq \pi/3, \\ 1_{\tilde{d}}, & \text{otherwise,} \end{cases}$$

and $f'_{i,j} \in U(A_n)$ for $(i, j) \in S^{(n)}$ with $i \geq 1$ by

$$f'_{i,j}(x) = \begin{cases} \exp(2i(t - \pi)e_{0,0}), & x = \iota_{i,j}^{(n)}(t), \quad \pi \leq t \leq 2\pi, \\ 1_{\tilde{d}}, & \text{otherwise.} \end{cases}$$

Remark that $[f'_{0,j}]_{K_1(A_n)} = [f_{0,j}^{(n)}] = g_{0,j}^{(n)}$, $j = 1, 2, \dots, r_0^{(n)}$, and $[f'_{i,j}]_{K_1(A_n)} = [f_{i,j}^{(n)}] = g_{i,j}^{(n)}$, $(i, j) \in S^{(n)}$ with $i \geq 1$. By the definitions of η , ξ , and ξ_c , it follows that $\eta_\nu(f'_{0,j}) \equiv 1_{3\tilde{d}}$, $\nu \neq j$, $(\xi \oplus \xi_c)(f'_{0,j}) \equiv 1_{(m-3r_0^{(n)})\tilde{d}}$, and

$$\eta_j(f'_{0,j})(x) = \begin{cases} \exp(2ite_{0,0}), & x = \tilde{t}_{0,j}^{(n)}(t), \quad 0 \leq t \leq \pi, \\ \exp(-2ite_{0,0}), & x = \iota_{i,j}^{(n)}(t), \quad \pi \leq t \leq 2\pi, \\ 1_{3\tilde{d}}, & \text{otherwise.} \end{cases}$$

Then we have that

$$[\varphi_{n,0}(f'_{0,j})]_{K_1(A(G_n, \tilde{p}_0', \tilde{p}_1'))} = [(\eta \oplus \xi \oplus \xi_c)(f'_{0,j})] = g_{0,j}^{(n)}.$$

By the definition of η it follows that $\eta_\nu(f'_{i,j}) \equiv 1_{3\tilde{d}}$, $\nu = 1, 2, \dots, r_0^{(n)}$, and by the definition of $\xi_\delta^{(n)}$ it follows that $\xi_\delta^{(n)}(f'_{i,j}) \equiv 1_{\tilde{d}}$ for $\delta \neq P_0 + P_1 + \sum_{\mu < i, p_\mu | G_{n,1}} r_\mu^{(n)} + j$ and $\xi_\delta(f'_{i,j}) = f'_{i,j}$ when $\delta = P_0 + P_1 + \sum r_\mu^{(n)} + j$. Then we have that

$$[\varphi_{n,0}(f'_{i,j})]_{K_1(A(G_n, \tilde{p}_0', \tilde{p}_1'))} = [(\eta \oplus \xi \oplus \xi_c)(f'_{i,j})] = g_{i,j}^{(n)}.$$

Hence we conclude that $(\varphi_{n,0}|_{A_n})_* = \text{id}_{G_n}$.

Proof of (1). Let $f \in C(\tilde{X}_n) \otimes M_{\tilde{d}}$. When $\varphi_{n,0}(f)^{-1}(\{0\}) \cap (\tilde{I}_{0,j'}^{(n)} \cup \tilde{I}_{1,j'}^{(n)})$ is ε -dense in $\tilde{I}_{0,j'}^{(n)} \cup \tilde{I}_{1,j'}^{(n)}$, by the definition of $D(t)$, we have that $f^{-1}(\{0\}) \cap (\tilde{I}_{0,j'}^{(n)} \cup \tilde{I}_{1,j'}^{(n)})$ is $1/3\varepsilon$ -dense in $\tilde{I}_{0,j'}^{(n)} \cup \tilde{I}_{1,j'}^{(n)}$. When $\varphi_{n,0}(f)^{-1}(\{0\}) \cap I_{i',j'}^{(n)}$ is ε -dense in $I_{i',j'}^{(n)}$ for any $(i', j') \in S^{(n)}$, by the definition of ξ_δ , we have that $f^{-1}(\{0\}) \cap I_{i,j}^{(n)}$ is $1/2\varepsilon$ -dense in $I_{i,j}^{(n)}$ for any $(i, j) \in S^{(n)}$. If $\varphi_{n,0}(f)^{-1}(\{0\}) \neq \emptyset$, then $f(c_n) = 0$ by ξ_c . Inductively, define relatively prime natural numbers $\tilde{p}_{0,m}, \tilde{p}_{1,m}$, and the unital embeddings $\varphi_{n,m} : A_{n,m-1} \hookrightarrow A_{n,m} = A(G_n, \tilde{p}_{0,m}, \tilde{p}_{1,m})$ by the same way as $\varphi_{n,0}$, then $\tilde{\varphi}_{n,0} = \varphi_{n,m} \circ \varphi_{n,m-1} \circ \cdots \circ \varphi_{n,0} : A_n \hookrightarrow A_{n,m}$ satisfies (1) (and (0)). Thus we may claim that $\varphi_{n,0}$ satisfies (1).

Proof of (2). Let x_0 and $y_0 \in \tilde{X}_n$ be such that $d(x_0, y_0) \leq \varepsilon$ and let $\hat{\eta}_{j,i} : \tilde{X}_n \rightarrow \tilde{X}_n$ for $i = -1, 0, 1$ be the continuous maps such that

$$\eta_j(f) = f \circ \hat{\eta}_{j,0} \oplus f \circ \hat{\eta}_{j,1} \oplus f \circ \hat{\eta}_{j,-1}.$$

By the definition of η_j , it follows that

$$d(\hat{\eta}_{j,i}(x), \hat{\eta}_{j,i}(y)) \leq 1/3d(x, y), \quad x, y \in \tilde{X}_n.$$

Then we have that $|\text{Tr}_{3r_0^{(n)}\tilde{d}}(\eta(f)(x_0) - \eta(f)(y_0))|$

$$\begin{aligned} &\leq 1/(3r_0^{(n)}) \sum_{j=1}^{r_0^{(n)}} \sum_{i=-1,0,1} |\text{Tr}_{\tilde{d}}(f \circ \hat{\eta}_{j,i}(x_0) - f \circ \hat{\eta}_{j,i}(y_0))| \\ &\leq \max\{|\text{Tr}(f(x) - f(y))|; d(x, y) \leq 1/3\varepsilon\}. \end{aligned}$$

Let $\hat{\xi}_\delta : \tilde{X}_n \rightarrow \tilde{X}_n$ be the continuous maps such that $\xi_\delta(f) = f \circ \hat{\xi}_\delta$. By the definition of ξ_δ , it follows that

$$d(\hat{\xi}_\delta(x), \hat{\xi}_\delta(y)) \leq 1/2d(x, y), \quad x, y \in \tilde{X}_n.$$

Then we have that $|\text{Tr}_{\kappa\tilde{d}}(\xi(f)(x_0) - \xi(f)(y_0))|$

$$\begin{aligned} &\leq 1/\kappa \sum_{\delta=1}^{\kappa} |\text{Tr}_{\tilde{d}}(f \circ \hat{\xi}_\delta(x_0) - f \circ \hat{\xi}_\delta(y_0))| \\ &\leq \max\{|\text{Tr}(f(x) - f(y))|; d(x, y) \leq 1/2\varepsilon\}. \end{aligned}$$

Combining the estimation for η and ξ , we have that

$$\begin{aligned} &|\text{Tr}_{\tilde{d}}(\varphi_{n,0}(f)(x_0) - \varphi_{n,0}(f)(y_0))| \\ &= |\text{Tr}((\eta \oplus \xi \oplus \xi_c)(f)(x_0) - (\eta \oplus \xi \oplus \xi_c)(f)(y_0))| \\ &\leq 3r_0^{(n)}/m |\text{Tr}_{3r_0^{(n)}\tilde{d}}(\eta(f)(x_0) - \eta(f)(y_0))| \\ &\quad + \kappa/m |\text{Tr}_{\kappa\tilde{d}}(\xi(f)(x_0) - \xi(f)(y_0))| \\ &\leq \max\{|\text{Tr}_{\tilde{d}}(f(x) - f(y))|; d(x, y) \leq 1/2\varepsilon\}. \end{aligned}$$

Thus we conclude (2).

Proof of (3). Let $h_k^{(n)} \in C(\tilde{X}_n)_+^1$, $n, k \in \mathbb{N}$ be such that $h_k^{(n)} \searrow \chi(E_n)$, $k \rightarrow \infty$. Then we have that $\lim_{k \rightarrow \infty} \text{Tr}_{\tilde{d}'}(\varphi_{n,0}(h_k^{(n)} \otimes 1_{\tilde{d}})(x))$

$$= \lim_k \text{Tr}((\eta \oplus \xi)(h_k^{(n)} \otimes 1_{\tilde{d}})(x) \oplus 0_{(m-(3r_0^{(n)}+\kappa))\tilde{d}}), \quad x \in \tilde{X}_n.$$

By the definition of η and ξ we have that

$$\begin{aligned} \lim_k \eta(h_k^{(n)} \otimes 1_{\tilde{d}}) &\leq \chi(E_n) \otimes 1_{3r_0^{(n)}\tilde{d}}, \\ \lim_k \xi(h_k^{(n)} \otimes 1_{\tilde{d}}) &\leq \chi(E_n) \otimes 1_{\kappa\tilde{d}}. \end{aligned}$$

Then, by $m \geq 2(3r_0^{(n)} + \kappa)$, we have that

$$\text{Tr}_{\tilde{d}'}(\varphi_{n,0}(\chi(E_n) \otimes 1_{\tilde{d}})(x)) \leq 1/2 \text{Tr}(\chi(E_n) \otimes 1_{\tilde{d}}(x)), \quad x \in \tilde{X}_n.$$

Theorem 3.2. *Let G be a countable abelian group. Then there exists a unital simple C^* -algebra A which has a unique tracial state, is expressed as the inductive limit C^* -algebra of an inductive sequence of dimension drop algebras, and satisfies $(K_0(A), [1_A]) \cong (\mathbb{Z}, 1)$ and $K_1(A) \cong G$.*

Proof. Let $(G_n)_{n \in \mathbb{N}}$ be an increasing sequence of finitely generated abelian groups and γ_n connecting maps such that $G = \varinjlim (G_n, \gamma_n)$. We may assume that γ_n is injective. By Proposition 2.2 and 3.1 we inductively obtain dimension drop algebras $A_{n,0}$ and A_n for G_n and unital embeddings $\psi_n : A_n \hookrightarrow A_{n+1,0}$ and $\varphi_{n+1,0} : A_{n+1,0} \hookrightarrow A_{n+1}$ such that $(\psi_n)_* \cong \gamma_n$, $(\varphi_{n+1,0})_* \cong \text{id}_{G_{n+1}}$, ψ_n satisfies the conditions (0,1) and (0,2) in Proposition 2.2, and $\varphi_{n+1,0}$ satisfies the conditions (1) and (2) in Proposition 3.1. Set $\varphi_n = \varphi_{n+1,0} \circ \psi_n$. Because of the conditions (0,1), (0,2), (1), and (2), we see that φ_n satisfies a similar conditions to (1) and (2) and satisfies $(\varphi_n)_* = \gamma_n$. Let A be the inductive limit C^* -algebra $\varinjlim (A_n, \varphi_n)$. It is not so hard to show that A is unital, projectionless, and satisfies $(K_0(A), [1_A]) \cong (\mathbb{Z}, 1)$ and $K_1(A) \cong G$.

By the condition (0,1) and (1) we shall show that A is simple. Let $\tilde{\varphi}_n$ be the canonical embedding from A_n into A , and let I be an ideal of A . Then it follows that

$$I = \overline{\bigcup \tilde{\varphi}_n(A_n) \cap I}.$$

Thus we obtain an ideal I_n of A_n such that $\tilde{\varphi}_n(A_n) \cap I = \tilde{\varphi}_n(I_n)$ and the compact subset $Y_n \subset X_n$ such that

$$I_n = \{f \in A_n; f|_{Y_n} = 0\}.$$

Assume that $A \neq I$, then there exists a subsequence $m_n \in \mathbb{N}$ such that $Y_{m_n} \neq \emptyset$, $n \in \mathbb{N}$. For any $n_0 \in \mathbb{N}$ and $f \in I_{n_0}$ we have that $\varphi_{m_n-1} \circ \varphi_{m_n-2} \circ \cdots \circ \varphi_{n_0}(f) \in$

I_{m_n} , for large $n \in \mathbb{N}$. By (0,1) and (1) of φ_n it follows that $f^{-1}(\{0\})$ is 2^{-m} -dense, where $m \rightarrow \infty$ ($m_n \rightarrow \infty$), and then $f = 0$, thus $I_{n_0} = \{0\}$ for any $n_0 \in \mathbb{N}$, which implies that $I = \{0\}$. Then we conclude that A is simple.

By the conditions (0,2) and (2), for $f \in A_n$ and $\varepsilon > 0$ we obtain $\delta > 0$ and $M \in \mathbb{N}$ such that if $m \geq M$ then

$$\begin{aligned} & \max\{|\operatorname{Tr}(\varphi_{m,n}(f)(x) - \varphi_{m,n}(f)(y))|; x, y \in X_{m+1}\} \\ & \leq \max\{|\operatorname{Tr}(\varphi_{m,n}(f)(x) - \varphi_{m,n}(f)(y))|; x, y \in \tilde{X}_{m+1}, d(x, y) \leq 4\pi\} \\ & \leq \max\{|\operatorname{Tr}(\varphi_{m-1,n}(f)(x) - \varphi_{m-1,n}(f)(y))|; x, y \in \tilde{X}_m, d(x, y) \leq 2\pi\} \\ & \vdots \\ & \leq \max\{|\operatorname{Tr}(f(x) - f(y))|; x, y \in \tilde{X}_n, d(x, y) \leq \delta\} \leq 1/2\varepsilon, \end{aligned}$$

where $\varphi_{m,n} = \varphi_m \circ \varphi_{m-1} \circ \cdots \circ \varphi_n$. Let $\tau_i \in T(A_{m+1})$, $i = 0, 1$, and let μ_{τ_i} be the probability measures on X_{m+1} such that

$$\tau_i \circ \varphi_{m,n}(f') = \int_{X_{m+1}} \operatorname{Tr}(\varphi_{m,n}(f')(x)) d\mu_{\tau_i}(x), \quad \text{for any } f' \in A_n.$$

By the above condition, it follows that $|\tau_0 \circ \varphi_{m,n}(f) - \tau_1 \circ \varphi_{m,n}(f)|$

$$\begin{aligned} & \leq \sum_{i=0,1} |\tau_i \circ \varphi_{m,n}(f) - \operatorname{Tr}(\varphi_{m,n}(f)(x_0))| \quad \text{for some } x_0 \in X_{m+1} \\ & \leq \varepsilon, \end{aligned}$$

for any $m \geq M$. Then for a sequence $\tau_m \in T(A_m)$, $m \in \mathbb{N}$ it follows that $(\tau_{m+1} \circ \varphi_{m,n}(f))_{m \in \mathbb{N}, m \geq n}$ is a Cauchy sequence and converges to a point which is independent of the choice of $\tau_m \in T(A_m)$. We define a tracial state τ on A by

$$\tau(\tilde{\varphi}_n(f)) = \lim_{m \rightarrow \infty} \tau_{m+1} \circ \varphi_{m,n}(f), \quad f \in A_n,$$

(remark that $\tau(\tilde{\varphi}_{l+1} \circ \varphi_{l,n}(f)) = \tau(\tilde{\varphi}_n(f))$). By the above claim for τ_0 and τ_1 , it follows that τ is unique. This completes the proof. $\blacksquare$

4 $\mathcal{Z}$ -stability of projectionless C^* -algebras

In this section, we prove that the projectionless C^* -algebra, that was constructed in Section 3, for a countable abelian group without free part absorbs the Jiang-Su algebra tensorially without passing through the classification theory of these algebras (Theorem 4.3). In order to prove $\mathcal{Z}$ -stability of the projectionless C^* -algebras for general countable abelian groups, we shall modify the constructions in Section 3 (Lemma 4.4 and Theorem 4.5).

The following lemma is a continuation of the proof of Proposition 3.1. Remark that if $G_n = G_{n,1}$ then we have that $\tilde{X}_n = X_n$. We set $\natural(0, i) = i$ for $i = 0, 1$, ($\natural(i) = 1 - i$, $d_{i,j}^{(n)} | \tilde{p}_{\natural(i,j)} \rangle$), $\tilde{d} = \tilde{p}_0 \tilde{p}_1$, $\tilde{d}' = P_0 P_1 \tilde{d}$, $A_n = A(G_n, \tilde{p}_0, \tilde{p}_1)$,

and $A_{n,0} = A(G_n, P_0 \tilde{p}_0, P_1 \tilde{p}_1)$. Let $\varphi_{n,0}$ be the unital embedding from A_n into $A_{n,0}$ that was defined in the proof of Proposition 3.1 and let A_n and $A_{n,0}$ be defined by unital embeddings $\{\epsilon_{i,j}^{(n)}\}$ and $\{\epsilon_{i,j}^{(n,0)}\}$.

Lemma 4.1. *Suppose that $G_n = G_{n,1}$. Then there exists a unital embedding ψ of $I(P_0, P_1)$ into $A_{n,0}$ satisfying that: if $f \in A_n$ and $\varepsilon > 0$ with $\|f(x) - f(y)\| < \varepsilon$ for any x and $y \in X_n$ then it follows that*

$$\|[\psi(g), \varphi_{n,0}(f)]\| < 2\varepsilon, \quad \text{for any } g \in I(P_0, P_1)^1.$$

Proof. We define C^* -subalgebras $A_{i,j}$, $(i, j) \in S^{(n)}$ of A_n by

$$A_{i,j} = \{f \in A_n; f(x) = f(a_{i,j}^{(n)}), x \in I_{i,j}^{(n)}\}.$$

By the definition of $\varphi_{n,0}$ in the proof of Proposition 3.1, it follows that

$$\begin{aligned} M_{\tilde{p}_i} &\cong \varphi_{n,0}(A_{0,i})(a_{0,i}^{(n)}) \subset \epsilon_{0,i}^{(n,0)}(M_{P_i \tilde{p}_i}), \quad i = 0, 1, \\ M_{\tilde{d}/d_{i,j}^{(n)}} &\cong \varphi_{n,0}(A_{i,j})(a_{i,j}^{(n)}) \subset \epsilon_{i,j}^{(n,0)}(M_{\tilde{d}'/d_{i,j}^{(n)}}), \quad (i, j) \in S^{(n)} \text{ with } i \geq 1 \end{aligned}$$

(where we denote by $A(x)$ the C^* -subalgebra $\{f(x); f \in A\}$ of $M_{\tilde{d}'}$ for a C^* -subalgebra A of $C(X_n) \otimes M_{\tilde{d}'}$ and $x \in X_n$.) then there exist unital embeddings $\delta_{0,i} : M_{P_i} \hookrightarrow \varphi_{n,0}(A_{0,i})(a_{0,i}^{(n)})' \cap \epsilon_{0,i}^{(n,0)}(M_{P_i \tilde{p}_i})$ for $i = 0, 1$, and $\delta_{i,j} : M_{P_{\natural(i,j)}} \hookrightarrow \varphi_{n,0}(A_{i,j})(a_{i,j}^{(n)})' \cap \epsilon_{i,j}^{(n,0)}(M_{\tilde{d}'/d_{i,j}^{(n)}})$, for $(i, j) \in S^{(n)}$ with $i \geq 1$.

We denote by $\psi' : C(X_n) \otimes M_{P_0 P_1} \hookrightarrow C(X_n) \otimes M_{\tilde{d}'}$ the natural unital embedding defined by $\psi'(f)(x) = 1_{\tilde{d}} \otimes f(x)$. By the definition of $\varphi_{n,0}$, we have that $\psi'(C(X_n) \otimes M_{P_0 P_1}) \subset \varphi_{n,0}(1_{C(X_n)} \otimes M_{\tilde{d}})'$. Set $I = I(P_0, P_1)$ and identify I with the C^* -subalgebra $\{f \in C(X_n) \otimes M_{P_0 P_1}; f(a_{0,0}^{(n)}) \in M_{P_0} \otimes 1_{P_1}, f(a_{0,1}^{(n)}) \in 1_{P_0} \otimes M_{P_1}, f(\iota_{i,j}^{(n)}(t)) = f(\iota_{0,\natural(i,j)}^{(n)}(t)) \text{ for } (i, j) \in S^{(n)}, i \geq 1, \text{ and } t \in [0, 2\pi]\}$. Since $\psi'(I)(a_{0,i}^{(n)}) \cong M_{P_i}$ and $\varphi_{n,0}(1_{C(\tilde{X}_n)} \otimes M_{\tilde{d}})(a_{0,i}^{(n)}) \supset \varphi_{n,0}(A_{0,i})(a_{0,i}^{(n)})$, we obtain a unitary $w_{0,i} \in U(C(X_n) \otimes M_{\tilde{d}'})$ such that $\text{Ad } w_{0,i} \circ \psi'(I)(a_{0,i}^{(n)}) = \delta_{0,i}(M_{P_i})$, $w_{0,i}(x) \in \varphi_{n,0}(A_{0,i})(a_{0,i}^{(n)})'$ for any $x \in I_{0,i}^{(n)}$, and $w_{0,i}(c_n) = 1$. Similarly, since $\psi'(I)(a_{i,j}^{(n)}) \cong M_{P_{\natural(i,j)}}$ and $\varphi_{n,0}(1_{C(\tilde{X}_n)} \otimes M_{\tilde{d}})(a_{i,j}^{(n)}) \supset \varphi_{n,0}(A_{i,j})(a_{i,j}^{(n)})$ for $(i, j) \in S^{(n)}$ with $i \geq 1$, we obtain a unitary $w_{i,j} \in U(C(X_n) \otimes M_{\tilde{d}'})$ such that $\text{Ad } w_{i,j} \circ \psi'(I)(a_{i,j}^{(n)}) = \delta_{i,j}(M_{P_{\natural(i,j)}})$, $w_{i,j}(x) \in \varphi_{n,0}(A_{i,j})(a_{i,j}^{(n)})'$ for any $x \in I_{i,j}^{(n)}$, and $w_{i,j}(c_n) = 1$. We define $w \in U(C(\tilde{X}_n) \otimes M_{\tilde{d}'})$ by

$$w(x) = w_{i,j}(x), \quad x \in I_{i,j}^{(n)}, \quad (i, j) \in S^{(n)},$$

and $\psi = \text{Ad } w \circ \psi'$, then it follows that $\psi(I) \subset A_{n,0}$.

Let $f \in A_n$ satisfy the condition in the lemma and let $f_{i,j} \in A_{i,j}$ for $(i, j) \in S^{(n)}$ be such that $f_{i,j}(a_{i,j}^{(n)}) = f(a_{i,j}^{(n)})$. By the definition of $\varphi_{n,0}$, we have that

$$\|\varphi_{n,0}(f_{i,j})(a_{i,j}^{(n)}) - \varphi_{n,0}(f)(x)\| = \left\| \bigoplus_{\delta=1}^m f_{i,j}(a_{i,j}^{(n)}) - f(x_\delta) \right\| < \varepsilon,$$

where $x_\delta \in X_n$, $m = P_0 P_1$. Hence it follows that for $g \in I(P_0, P_1)^1$ and $x \in I_{i,j}^{(n)}$, $(i, j) \in S^{(n)}$, $\psi(g)\varphi_{n,0}(f)(x) \approx_\varepsilon w_{i,j}\psi'(g)w_{i,j}^*(x) \varphi_{n,0}(f_{i,j})(a_{i,j}^{(n)}) = \varphi_{n,0}(f_{i,j})(a_{i,j}^{(n)}) w_{i,j}\psi'(g)w_{i,j}^*(x) \approx_\varepsilon \varphi_{n,0}(f)\psi(g)(x)$. Whence $\|[\psi(g), \varphi_{n,0}(f)]\| < 2\varepsilon$. $\blacksquare$

Lemma 4.2. *Let p and q be relatively prime numbers. If relatively prime numbers P_0 and P_1 satisfy $\min\{P_0, P_1\} \geq 2pq$, then there exists a unital embedding ψ of $I(p, q)$ into $I(P_0, P_1)$.*

Proof. Since $(p, q) = 1$ and $\min\{P_0, P_1\} > pq$, we obtain a, b, c , and $d \in \mathbb{Z}_+$ such that $P_0 = ap + bq$, $P_1 = cp + dq$, $b < p$, and $c < q$. Set $D = ad - bc$. Since $\min\{P_0, P_1\} \geq 2pq$ it follows that $D = \frac{1}{pq}(P_0 P_1 - P_1 bq - P_0 cp) > \frac{1}{pq}((P_0 - pq)(P_1 - pq) - (pq)^2) \geq 0$. Then we can separate $P_0 P_1$ as the following summation

$$P_0 P_1 = pcP_0 + qbP_1 + pqD.$$

We define a unital embedding $\psi : C(I_{0,0}^{(n)} \cup I_{0,1}^{(n)}) \otimes M_{pq} \hookrightarrow C(I_{0,0}^{(n)} \cup I_{0,1}^{(n)}) \otimes M_{P_0 P_1}$ by

$$\psi(f)(x) = V_0(f) \otimes 1_{cP_0} \oplus V_1(f) \otimes 1_{bP_1} \oplus f(x) \otimes 1_D,$$

where V_0 and V_1 are the irreducible representations of $I(p, q)$ at $a_{0,0}^{(n)}$ and $a_{0,1}^{(n)}$. Since $cP_0 + qD = aP_1$ and $bP_1 + pD = dP_0$, we obtain unital embeddings $\epsilon_{0,i} : M_{P_i} \hookrightarrow M_{P_0 P_1}$ such that $\psi(f)(a_{0,i}^{(n)}) \in \epsilon_{0,i}(M_{P_i})$, for any $f \in I(p, q)$ and $i = 0, 1$. Hence we have that $\psi(I(p, q)) \subset I(P_0, P_1)$, where $I(P_0, P_1)$ is defined by $\epsilon_{0,i}$. $\blacksquare$

Theorem 4.3. *The unital simple projectionless C^* -algebra, that is constructed in Theorem 3.2, for a countable abelian group without free part absorbs the Jiang-Su algebra $\mathcal{Z}$.*

Proof. By the definition of $\varphi_{n,0}$ in the proof of 3.1, for any $\varepsilon > 0$, $n_0 \in \mathbb{N}$, and any finite subset $F \subset A_{n_0}$ there exists a natural number $m \in \mathbb{N}$ such that $\max \{\|\varphi_{m,n}(f)(x) - \varphi_{m,n}(f)(y)\| : f \in F, x, y \in X_{m+1}\} < \varepsilon$, where $\varphi_{m,n} = \varphi_m \circ \varphi_{m-1} \circ \cdots \circ \varphi_n$ and $\varphi_n = \varphi_{n+1,0} \circ \psi_n : A_n \hookrightarrow A_{n+1}$ that was defined in the proof of Theorem 3.2. Applying Lemma 4.1 and Lemma 4.2, we have that

$$I(p, q) \subset_{\text{unital}} A' \cap A^\infty,$$

for any relatively prime numbers p and q . By Proposition 2.2 in [15], this implies that $A \otimes \mathcal{Z} \cong A$. $\blacksquare$

Lemma 4.4. *Let G_n be a finitely generated abelian group, p and q relatively prime numbers, $\tilde{p}_0$ and $\tilde{p}_1$ relatively prime numbers such that $d_{i,j}^{(n)} | \tilde{p}_0 \tilde{p}_1 = \tilde{d}$ for any $(i, j) \in S^{(n)}$ with $i \geq 1$, and A_n the dimension drop algebra for G_n , $\tilde{p}_0$, and $\tilde{p}_1$. Then there exist relatively prime numbers $\tilde{p}'_0$ and $\tilde{p}'_1$ such that $\tilde{p}_0 | \tilde{p}'_0$ and $\tilde{p}_1 | \tilde{p}'_1$, automorphisms $\{\epsilon_j^{(n,0)} \in \text{Aut}(M_{\tilde{d}'}); j = 1, 2, \dots, r_0^{(n)}\}$, where $\tilde{d}' = \tilde{p}_0 \tilde{p}'_1$,*

unital embeddings $\{\epsilon_{i,j}^{(n,0)} : M_{\tilde{d}'/d_{i,j}^{(n)}} \hookrightarrow M_{\tilde{d}'}; (i,j) \in S^{(n)}\}$, where $d_{0,i}^{(n)} = \tilde{p}_{\mathfrak{q}(i)}'$ for $i = 0, 1$, and a unital embedding $\varphi_{n,0} : C(\tilde{X}_n) \otimes M_{\tilde{d}} \hookrightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}'}$ satisfying the conditions (0) $\sim$ (3) in Proposition 3.1 and

- (4) there exist a unital embedding ψ of $I(p, q)$ into $A(G_n, \tilde{p}_0', \tilde{p}_1')$ satisfying that: if $f \in A_n$ and $\varepsilon > 0$ satisfy $\|f(x) - f(y)\| < \varepsilon$ for any x and $y \in X_n$ then it follows that

$$\|[\psi(g), \varphi_{n,0}(f)]\| < 2\varepsilon, \quad \text{for any } g \in I(p, q)^1.$$

Proof. Let $\alpha, \beta \in \mathbb{N}$ be such that $\alpha p - \beta q = 1$, $\alpha < q$, and $\beta < p$ and let P_i , $i = 0, 1$ be relatively prime numbers as in the proof of Proposition 3.1, satisfying moreover $P_0 P_1 > (\alpha p + \beta q) + \kappa$, $(P_i, pq) = 1$, and $\min\{P_0, P_1\} \geq 2pq$. Set $\tilde{p}_i' = P_i \tilde{p}_i$, $m = P_0 P_1$, and $\tilde{d}' = \tilde{p}_0' \tilde{p}_1'$.

Define a $*$ -homomorphism $\eta : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{(\alpha p + \beta q)\tilde{d}}$ by

$$\eta(f)(x) = \begin{cases} f(\tilde{t}_{i,j}^{(n)}(t)) \otimes 1_{\alpha p} \oplus f(\tilde{t}_{\mathfrak{q}(i),j}^{(n)}(t)) \otimes 1_{\beta q}, & x = \tilde{t}_{i,j}^{(n)}(t), t \in [0, \pi], \\ f(c_n) \otimes 1_{\alpha p + \beta q}, & \text{otherwise,} \end{cases}$$

define $\xi : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{\kappa \tilde{d}}$ and $\xi_c : C(\tilde{X}_n) \otimes M_{\tilde{d}} \rightarrow C(\tilde{X}_n) \otimes M_{(m - (\alpha p + \beta q + \kappa))\tilde{d}}$ by the same way as in the proof of Proposition 3.1, and define $\varphi_{n,0} : C(\tilde{X}_n) \otimes M_{\tilde{d}} \hookrightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}'}$ by $\varphi_{n,0} = \eta \oplus \xi \oplus \xi_c$. Remark that this $\varphi_{n,0}$ satisfies the condition (3). Let $v' \in U(C(\tilde{X}_n) \otimes M_{\tilde{d}})$ be such that $\text{Ad } v'(f)(b_{1,j}^{(n)}) = f(b_{0,j}^{(n)})$ for any $f \in A_n$ and $j = 1, 2, \dots, r_0^{(n)}$, and $v'(x) = 1_{\tilde{d}}$ for any $x \in \bigcup_{(i,j) \in S^{(n)}} I_{i,j}^{(n)}$, and set

$$v = v' \otimes 1_{\alpha p} \oplus v'^* \otimes 1_{\beta q} \oplus 1_{(m - (\alpha p + \beta q))\tilde{d}} \in U(C(\tilde{X}_n) \otimes M_{\tilde{d}'}).$$

Thus it follows that $\text{Ad } v \circ \varphi_{n,0}(f)(b_{1,j}^{(n)}) = \varphi_{n,0}(f)(b_{0,j}^{(n)})$ for any $f \in A_n$ and $j = 1, 2, \dots, r_0^{(n)}$. As in the proof of Proposition 3.1 for ξ , we obtain unital embeddings $\epsilon_{i,j}^{(n,0)}$, $(i,j) \in S^{(n)}$ such that $\varphi_{n,0}(f)(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n,0)}(M_{\tilde{d}'/d_{i,j}^{(n)}})$ for any $f \in A_n$ and $(i,j) \in S^{(n)}$. Let $A_{n,0}$ be the dimension drop algebra for G_n , $\tilde{p}_0'$, and $\tilde{p}_1'$ depending on the automorphisms $\{\text{Ad } v(b_{1,j}^{(n)})\}$ and the unital embeddings $\{\epsilon_{i,j}^{(n,0)}\}$, thus it follows that $\varphi_{n,0}(A_n) \subset A_{n,0}$. Since $\alpha p - \beta q = 1$, we have that $(\varphi_{n,0}|_{A_n})^* = \text{id}_{G_n}$.

As in the proof of Lemma 4.2 and $(P_i, pq) = 1$, we obtain a, b, c, d , and $D \in \mathbb{N}$ such that $D = ad - bc$ and $P_0 P_1 = pcP_0 + qbP_1 + pqD$. Set $I = I(p, q)$, and define a unital embedding $\psi' : I \hookrightarrow C(\tilde{X}_n) \otimes M_{\tilde{d}'}$ by $\psi'(g)(x)$

$$= 1_{\tilde{d}} \otimes (V_0(g) \otimes 1_{\alpha} \oplus V_1(g) \otimes 1_{\beta} \oplus g(x) \otimes 1_D \oplus V_0(g) \otimes 1_{cP_0 - \alpha} \oplus V_1(g) \otimes 1_{bP_1 - \beta}),$$

where V_0 and V_1 are the representations of I at $a_{0,0}^{(n)}$ and $a_{0,1}^{(n)}$, and we identify I with the C^* -subalgebra $\{f \in C(\tilde{X}_n) \otimes M_{pq}; f(a_{0,0}^{(n)}) \in M_p \otimes 1_q, f(a_{0,1}^{(n)}) \in$

$1_p \otimes M_q, f(\iota_{i,j}^{(n)}(t)) = f(\iota_{0,\mathfrak{h}(i,j)}^{(n)}(t))$ for $(i, j) \in S^{(n)}$ with $i \geq 1$ and $t \in [0, 2\pi]$, and $f(x) = f(c_n)$ for $x \in \tilde{I}_{i,j}^{(n)}$, $i = 0, 1$ and $j = 1, 2, \dots, r_0^{(n)}\}$. Then, for any $(i, j) \in S^{(n)}$, it follows that $\psi'(I)(a_{i,j}^{(n)})$ is contained unitally in a C^* -subalgebra of $\varphi_{n,0}(1_{C(\tilde{X}_n)} \otimes M_{\tilde{d}})(a_{i,j}^{(n)})' \cap M_{\tilde{d}'}$ that is isomorphic to $M_{P_{\mathfrak{h}(i,j)}}$. As in the proof of Lemma 4.1, we obtain a unitary $w \in U(C(\tilde{X}_n) \otimes M_{\tilde{d}'})$ such that $w(x) = 1_{\tilde{d}'}$ for $x \in \tilde{I}_{i,j}^{(n)}$, $\text{Ad } w \circ \psi'(g)(a_{i,j}^{(n)}) \in \epsilon_{i,j}^{(n,0)}(M_{\tilde{d}/d_{i,j}^{(n)}})$ for $g \in I$ and $(i, j) \in S^{(n)}$, and

$$\|[\text{Ad } w \circ \psi'(g)(x), \varphi_{n,0}(f)(x)]\| < 2\varepsilon,$$

for $g \in I^1$, $x \in \tilde{X}_n$, $f \in A_n$, and $\varepsilon > 0$ satisfying that $\|f(x) - f(y)\| < \varepsilon$ for any $x, y \in \tilde{X}_n$. Set $\psi = \text{Ad } w \circ \psi'$, then by the definitions of v and ψ' , it follows that

$$\text{Ad } v \circ \psi(g)(x) = \psi(g)(x), \quad x \in \tilde{X}_n,$$

which induces that $\psi(I) \subset A_{n,0}$.

In order to obtain the condition (1) and (2) for this $\varphi_{n,0}$, applying Proposition 3.1, we take the dimension drop algebra $A_{n,1} = A(G_n, \tilde{p}_0'', \tilde{p}_1'')$ and the unital embedding $\varphi_{n,1} : A_{n,0} \hookrightarrow A_{n,1}$, then $\varphi_{n,1} \circ \varphi_{n,0}$ satisfies (1) and (2). Hence we can construct $\varphi_{n,0}$ satisfies (0) $\sim$ (4). $\blacksquare$

Applying Proposition 2.2 and Lemma 4.4 in place of Proposition 3.1, we conclude the following theorem.

Theorem 4.5. *Let G be a countable abelian group. Then there exists a unital simple projectionless C^* -algebra A which has a unique tracial state, is expressed as the inductive limit C^* -algebra of an inductive sequence of dimension drop algebras, satisfies $(K_0(A), [1_A]) \cong (\mathbb{Z}, 1)$, $K_1(A) \cong G$, and absorbs the Jiang-Su algebra $\mathcal{Z}$.*

5 Certain aperiodic automorphism of projectionless C^* -algebras

In this section, we show the UHF-embeddability of the C^* -algebra in Theorem 3.2 or 4.5 (Proposition 5.2) and prove Theorem 1.2.

Let A be the unital simple projectionless C^* -algebra which is obtained in Theorem 3.2 or 4.5 for a countable abelian group G , τ the unique tracial state of A , (A_n, φ_n) the sequence of dimension drop algebras such that $A = \varinjlim (A_n, \varphi_n)$, where we have defined $\varphi_n : A_n \hookrightarrow A_{n+1}$ as the decomposition of unital embeddings such that

$$A_n = A(G_n, \tilde{p}_{0,n}, \tilde{p}_{1,n}) \xrightarrow{\psi_n} A(G_{n+1}, \tilde{p}'_{0,n+1}, \tilde{p}'_{1,n+1}) = A_{n+1,0} \xrightarrow{\varphi_{n+1,0}} A_{n+1}.$$

Set $\tilde{d}_n = \tilde{p}_{0,n}\tilde{p}_{1,n}$ and $\tilde{d}_{n,0} = \tilde{p}'_{0,n}\tilde{p}'_{1,n}$.

Let μ_n be the probability measure on X_n defined by $\tau \circ \tilde{\varphi}_n(f \otimes 1_{\tilde{d}_n}) = \int_{X_n} f(x) d\mu_n(x)$ for any $f \in C(X_n)$, and $E_n = \{a_{i,j}^{(n)}\} \cup \{\tau_j^{(n)}(z(\pi))\} \subset X_n$. We recall that ψ_n and $\varphi_{n+1,0}$ satisfy (0,3) and (3) in Proposition 2.2 and 3.1 or Lemma 4.4, then we have the following lemma.

Lemma 5.1. *For any $n \in \mathbb{N}$ we have that $\mu_n(E_n) = 0$.*

Proof. Let $h_k^{(n)} \in C(X_n)_+^1$ be such that $h_k^{(n)} \searrow \chi(E_n)$ (pointwise). By the conditions (0,3) and (3) in Proposition 2.2 and 3.1 or Lemma 4.4 we have that

$$\begin{aligned} \mu_n(E_n) &= \lim_{k \rightarrow \infty} \tau \circ \tilde{\varphi}_n(h_k^{(n)} \otimes 1_{\tilde{d}}) = \lim_k \tau \circ \tilde{\varphi}_{n+1} \circ \varphi_{n+1,0} \circ \psi_n(h_k^{(n)} \otimes 1_{\tilde{d}}), \\ &\leq \lim_k \tau \circ \tilde{\varphi}_{n+1} \circ \varphi_{n+1,0}(h_k^{(n+1)} \otimes 1_{\tilde{d}_{n+1,0}}) \\ &\leq \lim \int_{X_{n+1}} \text{Tr}_{\tilde{d}_{n+1}}(\varphi_{n+1,0}(h_k^{(n+1)} \otimes 1_{\tilde{d}_{n+1,0}})(x)) d\mu_{n+1}(x) \\ &\leq 1/2 \lim \int_{X_{n+1}} \text{Tr}_{\tilde{d}_{n+1}}(h_k^{(n+1)} \otimes 1_{\tilde{d}_{n+1}}(x)) d\mu_{n+1}(x) \\ &= 1/2 \mu_{n+1}(E_{n+1}). \end{aligned}$$

Hence we have that $\mu_n(E_n) = 0$. ■

Let (π_τ, H_τ) be the GNS-representation of A associated with τ . The following Proposition is a generalization of Proposition 2.3 in [13].

Proposition 5.2. *There exists a UHF C^* -subalgebra B of $\pi_\tau(A)''$ and an increasing sequence $(B_n)_{n \in \mathbb{N}}$ of matrix C^* -subalgebras of B such that $\pi_\tau(A) \subset B$, $1_{B_n} = 1_B$, $B = \overline{\bigcup B_n}^{\|\cdot\|}$, and*

$$(B'_n \cap \pi_\tau(A))'' = B'_n \cap \pi_\tau(A)''. \quad (5.1)$$

Proof. Let C_n denote the C^* -algebra $C(\tilde{X}_n) \otimes M_{\tilde{d}_n}$, where $\tilde{d}_n$ is such that $A_n \subset C(\tilde{X}_n) \otimes M_{\tilde{d}_n}$ and let $f_k \in C(X_n)_+^1$, $k \in \mathbb{N}$ be such that $f_k \nearrow \chi(X_n \setminus E_n)$, $k \rightarrow \infty$ (pointwise), and $f_k(E_n) = \{0\}$ and define a map $\Psi_n : C_n \rightarrow \pi_\tau(A)''$ by

$$\Psi_n(g) = \lim_{k \rightarrow \infty} \pi_\tau \circ \tilde{\varphi}_n((f_k \otimes 1_{\tilde{d}})g), \quad g \in C_n,$$

where the limit is taken in the strong operator topology. From Lemma 5.1 and a similar argument to the proof of Proposition 2.3 in [13], it follows that Ψ_n is independent of the choice of f_k , is an injective $*$ -homomorphism, $\Psi_n|_{A_n} = \pi_\tau \circ \tilde{\varphi}_n$ and satisfies $\Psi_n = \Psi_{n+1} \circ \varphi_n$ on C_n . Then we have that

$$\pi_\tau(A) \subset \overline{\bigcup_n \Psi_n(C_n)}^{\|\cdot\|} \subset \pi_\tau(A)''. \quad (5.2)$$

Set $B = \overline{\bigcup_n \Psi_n(C_n)}^{\|\cdot\|}$ and define a sequence of finite dimensional C^* -subalgebras B_n of B by

$$B_n = \Psi_n(1_{C(\tilde{X}_n)} \otimes M_{\tilde{d}_n}), \quad n \in \mathbb{N}.$$

By the definition of $\varphi_{n+1,0}$ and ψ_n , for $f \in C_n$ there exists a large natural number m such that $\varphi_m \circ \varphi_{m-1} \circ \cdots \circ \varphi_n(f)$ is almost contained in $1_{C(\tilde{X}_{m+1})} \otimes M_{d_{m+1}}$, then we have that $\Psi_n(C_n) \subset \overline{\bigcup B_n}$, hence $B = \overline{\bigcup B_n}$ is a UHF-algebra.

We shall show $(B'_n \cap \pi_\tau(A))'' \supset B'_n \cap \pi_\tau(A)''$. From Lemma 5.1 we have that

$$\Psi_N((1 - f_k) \otimes 1_{\tilde{d}_N}) = \pi_\tau \circ \tilde{\varphi}_N((1 - f_k) \otimes 1_{\tilde{d}_N}) \searrow 0, \quad k \rightarrow \infty \text{ (strongly),}$$

for any $N \in \mathbb{N}$. Let $g \in B'_n \cap \pi_\tau(A)''$. We may assume that $g \in B'_n \cap \Psi_N(C_N)$, $N > n$. Set $B_{n,0} = \varphi_{N-1,n}(1 \otimes M_{\tilde{d}_n}) \subset C_N$. We obtain $g_0 \in C_N \cap B'_{n,0}$ such that $g = \Psi_N(g_0)$. Since $g_0 \cdot (f_k \otimes 1_{\tilde{d}_N}) \in A_N \cap B'_{n,0}$ and $\Psi_N(g_0 \cdot (f_k \otimes 1_{\tilde{d}_N})) \rightarrow \Psi_N(g_0)$ (strongly), we have that $g \in (\pi_\tau(A) \cap B'_n)''$. $\blacksquare$

Remark 5.3. By the UHF-embeddability in the above proposition and by the same argument as in the proof of Theorem 2.6 in [13], we can prove 1.1 for the projectionless C^* -algebras that are constructed in this paper.

The following lemma is an adaptation of Theorem 3.12.14 in [11].

Lemma 5.4. *Suppose that a sequence $(F_n)_{n \in \mathbb{N}}$ of projections in $\pi_\tau(A)''$ satisfies that $[F_n, x] \rightarrow 0$ (strongly) for any $x \in \pi_\tau(A)''$. Then there exists a central sequence $(f_n)_n$ of positive elements in A^1 (in the sense that*

$$\|[f_n, a]\| \rightarrow 0, \text{ for any } a \in A,)$$

such that

$$F_n - \pi_\tau(f_n) \rightarrow 0 \quad (\text{strongly}).$$

We recall that

$$\text{WInn}(A) = \{\alpha \in \text{Aut}(A) : \pi_\tau \circ \alpha = \text{Ad } W \circ \pi_\tau, \quad W \in U(\pi_\tau(A)'')\}.$$

Mimicking the proof of Lemma 4.4 and Theorem 4.5 in [7], we obtain the following proposition.

Proposition 5.5. *Let A be a unital simple projectionless C^* -algebra with a unique tracial state τ and let $\alpha \in \text{Aut}(A)$. If there exists a central sequence $(f_n)_{n \in \mathbb{N}} \in (A_\infty)_+^1$ such that*

$$(\alpha(f_n))_n \cdot (f_n)_n = 0, \quad \tau(1 - (f_n + \alpha(f_n))) \rightarrow 0, \quad n \rightarrow \infty,$$

then $\alpha \notin \text{WInn}(A)$.

Proof. Assume that there exists $V \in U(\pi_\tau(A)'')$ such that $\pi_\tau \circ \alpha = \text{Ad } V \circ \pi_\tau$. Let $\rho : A \times_\alpha \mathbb{Z} \rightarrow \pi_\tau(A)''$ be a representation of $A \times_\alpha \mathbb{Z}$ determined by $\rho(a) = \pi_\tau(a)$ for $a \in A$ and $\rho(u_\alpha) = V$, where u_α is the canonical unitary with $u_\alpha a = \alpha(a)u_\alpha$, $a \in A$. Define a tracial state ϕ of $A \times_\alpha \mathbb{Z}$ by $\phi(x) = (\rho(x)\Omega_\tau | \Omega_\tau)$, where Ω_τ is the cyclic vector obtained by the GNS-representation associated with τ . By $V \in U(\pi_\tau(A)'')$ there are $a_n \in A$ such that $\pi_\tau(a_n) \rightarrow V^*$ (strongly). Thus we have $\phi(a_n u_\alpha) \rightarrow 1$. However we shall obtain $\phi(a u_\alpha) = 0$ for any $a \in A$ by the condition of α and the following argument.

Set $a \in A^1$ and let $(f_n) \in (A_\infty)_+^1$ be as in the Proposition, then it follows that

$$\begin{aligned}\phi(au_\alpha) &= \lim_{n \rightarrow \infty} \phi((f_n + \alpha(f_n))au_\alpha) + \phi((1 - (f_n + \alpha(f_n)))au_\alpha) \\ &= \lim_{n \rightarrow \infty} \phi((f_n + \alpha(f_n))au_\alpha).\end{aligned}$$

Since $(f_n)_n$ is a central sequence we have that

$$\lim_{n \rightarrow \infty} \phi(f_n au_\alpha) = \lim_{n \rightarrow \infty} \phi(f_n^{1/2} \alpha(f_n)^{1/2} au_\alpha) = 0,$$

which implies that $\phi(au_\alpha) = 0$. ■

The following proof of Theorem 1.2 is based on the proof of Theorem 4.5 in [7].

Proof of Theorem 1.2. Suppose that $\alpha \in \text{Aut}(A)$ has a central sequence $(f'_n)_n \in (A_\infty)_+^1$ which satisfies the condition in the theorem for $2k$ in place of k . Set $f_n = \sum_{j=0}^{k-1} \alpha^j(f'_n)$. Thus $(f_n)_n \in (A_\infty)_+^1$ satisfies $(f_n) \cdot \alpha^k((f_n)) = 0$, and $\tau(1_A - (f_n + \alpha^k(f_n))) \rightarrow 0$. From Proposition 5.5, it follows that $\alpha^k \notin \text{WInn}(A)$. Then we have that $[\alpha]$ is aperiodic in $\text{Aut}(A)/\text{WInn}(A)$.

Suppose that $[\alpha] \in \text{Aut}(A)/\text{WInn}(A)$ is aperiodic. Let $\bar{\alpha}$ be the weak extension of $\pi_\tau \circ \alpha \circ \pi_\tau^{-1}$ to an automorphism of $\pi_\tau(A)''$. Let $k \in \mathbb{N}$, $\varepsilon_n > 0$, $n \in \mathbb{N}$, be such that $\varepsilon_n \searrow 0$, and F_n , $n \in \mathbb{N}$, be finite subsets of A^1 such that $\overline{(\bigcup F_n)}^{\|\cdot\|} = A^1$. By the classification theory for aperiodic automorphisms on the injective type II_1 factor due to Connes [1], there exists a central sequence $(E_m) \in (\pi_\tau(A)'')_\infty$ in the sense of the strong topology of projections such that

$$E_m + \bar{\alpha}(E_m) + \cdots + \bar{\alpha}^{k-1}(E_m) \rightarrow 1 \text{ (strongly)}.$$

By Lemma 5.4 we obtain a central sequence $(f'_m)_m \in A_\infty$ in the sense of the norm such that $f'_m \in A_+^1$ and $E_m - \pi_\tau(f'_m) \rightarrow 0$ (strongly). Then we have that

$$\tau(1 - \sum_{j=0}^{k-1} \bar{\alpha}^j \circ \pi_\tau(f'_m)) \rightarrow 0, \quad m \rightarrow \infty. \text{ Set}$$

$$g_m = (f'_m)^{1/2}(\alpha(f'_m) + \alpha^2(f'_m) + \cdots + \alpha^{k-1}(f'_m))(f'_m)^{1/2}, \quad m \in \mathbb{N},$$

then we have $\|\pi_\tau(g_m) - E_m(\bar{\alpha}(E_m) + \bar{\alpha}^2(E_m) + \cdots + \bar{\alpha}^{k-1}(E_m))E_m\|_2$

$$\begin{aligned}&\leq \|(\pi_\tau(f'_m)^{1/2} - E_m)(\sum_{j=1}^{k-1} \bar{\alpha}^j(\pi_\tau(f'_m)))\pi_\tau(f'_m)^{1/2}\|_2 \\ &+ \|E_m(\sum_{j=1}^{k-1} \bar{\alpha}^j(\pi_\tau(f'_m) - E_m))\pi_\tau(f'_m)^{1/2}\|_2 \\ &+ \|E_m(\sum_{j=1}^{k-1} \bar{\alpha}^j(E_m))(\pi_\tau(f'_m)^{1/2} - E_m)\|_2 \\ &\leq 2k\|\pi_\tau(f'_m)^{1/2} - E_m\|_2 + k\|\pi_\tau(f'_m) - E_m\|_2 \rightarrow 0\end{aligned}$$

and by $\|E_m(\bar{\alpha}(E_m) + \bar{\alpha}^2(E_m) + \cdots + \bar{\alpha}^{k-1}(E_m))E_m\|_2 \rightarrow 0$, we have that $\|\pi_\tau(g_m)\|_2 \rightarrow 0$.

Define $f_\varepsilon \in C([0, \infty))_+^1$ by

$$f_\varepsilon(t) = \begin{cases} (2\varepsilon)^{-2}t, & 0 \leq t \leq 4\varepsilon^2, \\ 1 & 4\varepsilon^2 < t, \end{cases}$$

and let

$$f_{m,n} = (f'_m)^{1/2} \cdot (1_A - f_{\varepsilon_n}(g_m)) \cdot (f'_m)^{1/2} \in A_+^1, \quad m, n \in \mathbb{N}.$$

Remark that $0 \leq f_{m,n} \leq f'_m$ and $(f_{m,n})_m$ is a central sequence in the norm sense for any $n \in \mathbb{N}$ (i.e., $\|[f_{m,n}, a]\| \rightarrow 0$, $m \rightarrow \infty$, $a \in A$). Then for any $j = 1, 2, \dots, k-1$ and $m, n \in \mathbb{N}$ we have that $\|f_{m,n}\alpha^j(f_{m,n})\|^2$

$$\begin{aligned} &\leq \|f_{m,n}\alpha^j(f_{m,n})^{1/2}\|^2 = \|f_{m,n}\alpha^j(f_{m,n})f_{m,n}\| \\ &\leq \|f_{m,n}(\alpha(f'_m) + \alpha^2(f'_m) + \dots + \alpha^{k-1}(f'_m))f_{m,n}\| \\ &\leq \|(f'_m)^{1/2}(\alpha(f'_m) + \dots + \alpha^{k-1}(f'_m))(f'_m)^{1/2}(1_A - f_{\varepsilon_n}(g_m))(f'_m)^{1/2}\| \\ &\leq \|g_m - g_m \cdot f_{\varepsilon_n}(g_m)\| = \|g_m \cdot (1 - f_{\varepsilon_n})(g_m)\| = \varepsilon_n^2. \end{aligned}$$

Since $\|\pi_\tau(g_m)\|_2 \rightarrow 0$ and $(f_{m,n})_m \in A_\infty$, for $\varepsilon_n > 0$ and F_n we obtain a sub-sequence $m_n \in \mathbb{N}$ such that

$$\|f_{m_n,n} - f'_{m_n}\|_2 \leq \|f_{\varepsilon_n}(g_{m_n})\|_2 < \varepsilon_n,$$

$$\|[f_{m_n,n}, x]\| < \varepsilon_n, \quad x \in F_n.$$

Let $f_n = f_{m_n,n} \in (A)_+^1$. Thus $(f_n)_n$ is a central sequence in the norm sense which satisfies

$$(f_n)_n(\alpha^j(f_n))_n = 0, \quad j = 1, 2, \dots, k-1.$$

Since $\tau(1 - \sum_{j=0}^{k-1} \overline{\alpha}^j \circ \pi_\tau(f'_{m_n})) \rightarrow 0$, ($n \rightarrow \infty$), we have $\tau(1 - \sum_{j=0}^{k-1} \alpha^j(f_n)) \rightarrow 0$.

This completes the proof. ■

Acknowledgment. The author is grateful to Professor Akitaka Kishimoto for his support and valuable comments.

References

- [1] A. Connes. *Outer conjugacy class of automorphisms of factors*, Ann. Scient. Ec. Norm. Sup. (4) **8** (1975), 383-420.
- [2] M. Dadarlat and W. Winter *Trivialization of $C(X)$ -algebras with strongly self-absorbing fibres*, Bull. Soc. Math. France 136 (2008), no. 4, 575-606.
- [3] M. Dadarlat, N. C. Phillips, and A. S. Toms *A direct proof of $\mathcal{Z}$ -stability for AH algebras of bounded topological dimension*, arXiv:0806.2855 17 Jun 2008.

- [4] G. A. Elliott *An invariant for simple C^* -algebras*, Canadian Mathematical Society. 1945-1995 3 (1996), 61-90.
- [5] G. A. Elliott and A. S. Toms *Regularity properties in the classification program for separable amenable C^* -algebras*, Bull. Amer. Math. Soc. 45 (2008), 229-245.
- [6] X. Jiang and H. Su. *On a simple unital projectionless C^* -algebra*, Amer. J. Math. 121 (1999), no. 2, 359–413.
- [7] A. Kishimoto. *The Rohlin property for shifts on UHF algebras and automorphisms of Cuntz algebras*, J. Funct. Anal. 140 (1996), no. 1, 100–123.
- [8] L. Li. *Classification of simple C^* -algebras: inductive limits of matrix algebras over trees*, Mem. Amer. Math. Soc. 127 (1997), no. 605, vii+123 pp.
- [9] H. Lin. *Inductive Limits of Subhomogeneous C^* -algebras with Hausdorff Spectrum*, arXiv:0809.5273 30 Sep 2008.
- [10] J. Mygind. *Classification of certain simple C^* -algebras with torsion in K_1* , Canad. J. Math. 53 (2001), no. 6, 1223-1308.
- [11] G. K. Pedersen. “ C^* -algebras and their automorphism groups.” London Mathematical Society Monographs, 14. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1979.
- [12] M. Rørdam and W. Winter. *The Jiang-Su algebra revisited*, arXiv:0801.2259
- [13] Y. Sato. *Certain aperiodic automorphisms of unital simple projectionless C^* -algebras*, submitted.
- [14] K. Thomsen *Limits of certain subhomogeneous C^* -algebras*, Mem. Soc. Math. Fr. (N.S.) No. 71 (1997), vi+125 pp. (1998).
- [15] A. S. Toms and W. Winter. *$\mathbb{Z}$ -stable ASH algebras*, arXiv:math/0508218
- [16] W. Winter. *Localizing the Elliott conjecture at strongly self-absorbing C^* -algebras -with an appendix by Huaxin Lin-*, arXiv:0708.0283 11 Sep 2007.